

# CALCULATED SEDIMENTATION RATE IN THE KRKA RIVER ESTUARY USING VERTICAL DISTRIBUTION OF $^{137}\text{Cs}$

Neven Cukrov <sup>1</sup>\*, Delko Barišić <sup>1</sup> and Mladen Juračić <sup>2</sup>

<sup>1</sup> Rudjer Bošković Institute, Bijenička 54, 10000 Zagreb, Croatia. - ncukrov@irb.hr

<sup>2</sup> Faculty of Science, University of Zagreb, Horvatovac 102 a, 10000 Zagreb, Croatia

## Abstract

Sedimentation rate in the Krka River estuary was calculated from vertical  $^{137}\text{Cs}$  distribution in sediments. Calculated sedimentation rate was 2 mm/a upstream of the Prokljan Lake, 4-5 mm/a at the Guduča River mouth in Prokljan Lake and 3-4 mm/a in other parts of the lake. In the lower part of estuary, sedimentation rate was very small, less than 1 mm with exceptions in deepest parts of estuary in front of city Šibenik where sedimentation rate was approx. 3 mm/a.

**Keywords :** Adriatic Sea, Estuaries, Radionuclides, Sedimentation.

The Krka River estuary was formed during the Holocene transgression. Located between Skradinski Buk waterfalls (calc-tufa barrier) through the Prokljan Lake to the St. Nikola fortress, the estuary has a total length of 22 km (Fig 1).




Fig. 1. Krka River estuary, with sampling locations and four different sedimentation areas recognized. Inset: Characteristic  $^{137}\text{Cs}$  activities in sediment profiles at investigated locations.

The hydrogeologic drainage area of the Krka River is approximately 2427 km<sup>2</sup>. The estuary bottom gradually deepens from 2 m, below the waterfalls, to 42 m in front of Fort St. Nikola. It is typical karstic highly stratified estuary with fresh/brackish surface layer flowing seawards and bottom seawater counter-current moving upward. Input of terrigenous clastic material into the Krka river estuary is relatively small [1], with main input of the particulate material in the Krka River estuary via small Guduča River (Fig 1) inflowing into the Prokljan Lake [1], [2]. The Krka River carries larger quantities of fresh-water (on average 55 m<sup>3</sup>/s) than the Guduča River (on average less than 1 m<sup>3</sup>/s). However, a number of calc tufa barriers along the Krka River, upstream of the town of Skradin, significantly reduce suspended material transport [1], [2], [3].

The bottom sediment samples were collected by a scuba diver using hand-driven plexyglas corers, at 6 locations in the Krka River Estuary. Prior to the gamma-spectrometry measurements, the sediment samples were dried

at 106°C during 24 hours, counted in a special vessels, sealed and stored for at least 4 weeks in order to allow a radioactivity disintegration of a gaseous  $^{222}\text{Rn}$ . The samples were counted on an HPGe detector with an 8192 channel analyzer. The system was calibrated using the standards supplied by Amersham International, IAEA-306 and IAEA-314. Sedimentation rate was calculated using the vertical distribution of  $^{137}\text{Cs}$ , (Fig 1 inset).

Sedimentation rate in the estuary was found to be very low (0,27 mm/a) based on  $^{14}\text{C}$  measurements [1]. First attempts using  $^{137}\text{Cs}$  distribution indicated sedimentation rate higher than 2 mm/a ([4]). According to the prevailing origin of accumulated material and sedimentation rates, four main sedimentation areas in the Krka River Estuary were recognized [5]. Anthropogenic radionuclides may be released in the environment owing to nuclear explosions, testing of nuclear weapons and discharge of effluents from nuclear facilities. Anthropogenic radionuclide  $^{137}\text{Cs}$  with half life of 30,18 a first time entered in the environment as a product of atomic explosions in 1945. Since the Second World War, more than 2400 nuclear weapon experiments have been conducted worldwide [6] and as a result  $^{137}\text{Cs}$  spread around the world and become good sediment marker.

In the III sedimentation area where most of the deposited material is carried by the Krka River, calculated sedimentation rate was 2 mm/a, while in the Prokljan Lake where recent sediment consists mainly of particles of terrigenous flysch transported by the Guduča River (IV sedimentation area) was 3-4 mm/a. The highest sedimentation rate was measured at the Guduča River mouth (4-5 mm/a). In the II sedimentation area deposited material is mixture of particles from two main sources: marine carbonates and terrigenous flysch, and the sedimentation rate is less than 2 mm. In the I sedimentation area where carbonate marine sedimentation prevails, sedimentation rate is probably less than 1 mm/a, with exceptions in the deepest parts of estuary where sedimentation rate is approximately 3 mm/a. This deepest area is located at the center of the lower part of the estuary and acts as a kind of sediment trap (Fig 1).

## References

- 1 - Juračić M. and Prohić E. 1991. Mineralogy, Sources of particles, and sedimentation in the Krka River estuary (Croatia). *Geološki Vjesnik* 44: 195-200.
- 2 - Cukrov N., Barišić D. and Branica M. 2004. Content of the  $^{40}\text{K}$ ,  $^{232}\text{Th}$ ,  $^{226}\text{Ra}$ ,  $^{238}\text{U}$  and  $^{137}\text{Cs}$  in the recent sediments of the Krka river estuary. XXXVIIth *CIESM. Congress Proc.*, 37: 20.
- 3 - Prohić E. and Juračić M. 1989. Heavy Metals in sediment - Problems Concerning Determination of Anthropogenic Influence. Study in the Krka River Estuary, Eastern Adriatic, *Environ. Geol. Water Sci.* 13 (2): 145-151.
- 4 - Cukrov N., Barišić D., Branica M. and Kwokal Ž. 2003. Preliminary results of Uranium and Radium content in Krka River Estuary. Abstracts book, 22nd IAS Meeting of Sedimentology: 36.
- 5 - Cukrov N. and Barišić D. 2006. Spatial Distribution of  $^{40}\text{K}$  and  $^{232}\text{Th}$  in Recent Sediments of the Krka. *Croatia Chemica Acta* 79 (1): 115-118.
- 6 - Norris R.S. and Arkin W.M. 1998. Known nuclear tests worldwide, 1945-98. *Bulletin of the Atomic Scientists*, 54 (06): 65-67.