

The Flora of Zečevo (Hvar Archipelago, Croatia)

Mario SKELIN¹

Ivica LJUBIĆ² (✉)

Ivan SKELIN¹

Ivana VITASOVIĆ KOSIĆ²

Sandro BOGDANOVIĆ²

Summary

The islet of Zečevo belongs to Hvar Archipelago and it is situated two kilometres north-east of Vrboska on the island of Hvar (Croatia). The area of islet is protected as a Significant landscape. During vegetation periods from 2006 to 2010 a total of 216 vascular plant taxa were recorded. From those, 12 taxa were recorded for Hvar Archipelago for the first time. The taxa were classified within 65 families and 161 genera. The most abundant families are legumes (*Fabaceae* 13%), daisies (*Asteraceae* s.l. 9.2%), grasses (*Poaceae* 8.3%) and mints (*Lamiaceae* 5.1%). According to the analysis of life forms the largest portion of therophytes (45.4%) and hemicryptophytes (21.3%) were found. Phytogeographical analysis reveals predominance of the Mediterranean floral element (63.4%), followed by widespread plants (12.5%) and Euro-Asian floral element (9.7%). Five endemics and five endangered species were recorded, among them three are vulnerable (VU): *Ophrys bertolonii* Moretti, *O. sphegodes* Mill., *Salsola kali* L. and two are endangered (EN): *Glaucium flavum* Crantz and *Carex extensa* Gooden. Three rare plant species of Croatian flora: *Ophrys liburnica* Devillers et Devillers-Tersch, *Scaligeria cretica* (Mill.) Boiss. and *Narcissus serotinus* L. were found.

Key words

Zečevo, islet of Hvar, vascular flora, plant diversity, Croatia

¹ Mala banda bb, 21465 Jelsa, Island of Hvar, Croatia

² University of Zagreb, Faculty of Agriculture, Department of Agricultural Botany, Svetosimunska 25, HR-10000 Zagreb, Croatia

✉ e-mail: iljubicic@agr.hr

Received: November 26, 2013 | Accepted: April 3, 2014

Introduction

The islet of Zečevo has a status of Significant landscape since 1968 (Gabelica et al. 2011). The islet is uninhabited, which excludes major anthropogenic influence on composition and dynamics of flora. Except diverse floral composition the islet is inhabited by various animals, primarily with the high population of rabbits (in Croatian "zec") from which this islet got the name after.

First floristic research of the island of Hvar was conducted in 1818 when Franjo Portenschlag-Ledermayer visited the island and explored its flora. Since then, according to Trinajstić (1993) many individual researches of flora were carried out for example: Visiani in 1842-1852, Biasoletto in 1841, Alschinger in 1861, Pittoni in 1869, Huter in 1870 and Studnička in 1890. Trinajstić (1993) incorporated all previous floristic data and a total of 1095 plant taxa was published in "Flora of the island of Hvar".

Despite numerous floristic studies from central and southern Dalmatian islands (e.g. Trinajstić, 1993, 1998; Jasprica and Kovačić, 1997; Vladović and Jukić, 1997; Kovačić et al., 2000; Vladović et al., 2001, 2002; Bogdanović and Mitić, 2003; Bogdanović et al., 2004; Milović, 2004; Jasprica et al., 2006) the flora of Zečevo islet was never investigated before. The aim of this study was to investigate floristic diversity and to compare it with other smaller islets of the Adriatic area.

Materials and methods

Study area

The islet of Zečevo is situated two kilometres north-east of town Vrboska on the island of Hvar in central Dalmatia (Fig. 1). Surface of Zečevo islet is approximately 0.11 km², while the coastal length is 1539 m (Duplančić Leder et al., 2004). On the islet vegetation of *Pinus halepensis* L. woods and macchias under the influence of typical Mediterranean climate is developed (Mihovilović, 1995). Field work on Zečevo islet was carried out within four vegetation seasons between 2006 and 2010 in different periods.




Figure 1. The geographical position of Zečevo islet

Materials and methods

The standard methods for floristic mapping of the collection and preparation of floristic materials and determination were used according to Nikolić (2006). Nomenclature has been adjusted according to the Flora Croatica Database (Nikolić, 2013a). Families, genera, species and subspecies are listed in alphabetical order and organized according to higher systematic units (Nikolić, 2013b).

The analysis of life forms was made according to Raunkiaer (1934), Horvat (1949) and Pignatti (1982). In the floristic list, life forms are marked with the following abbreviations: *Chamaephyta* (Ch), *Geophyta* (G), *Hemicryptophyta* (H), *Phanerophyta* (P) and *Therophyta* (T). The analysis of floral elements was made according to Horvatić (1963). For corresponding floral element the following abbreviations are used: (1) Mediterranean: circum Mediterranean plants – cm, east Mediterranean plants – em, Illyrian Adriatic plants – ila, Mediterranean Atlantic plants – ma, European Mediterranean plants – eum, Mediterranean Pontic plants – mp, (2) South European: south European Mediterranean plants – sem, south European Pontic plants – sep, (3) Southeast European – se, (4) European – eu, (5) Euro-Asian – ea, (6) Widespread plants – wp and (7) Alien plants – al.

Endemic (e) and invasive species (Inv) are determined according to the Flora Croatica Database (Nikolić, 2013a) and according to Boršić et al. (2008). Endangered plant species, following the abbreviations: critically endangered (CR), endangered (EN), vulnerable (VU), least concern (LC), near threatened (NT), and data deficient (DD) were determined according to Nikolić and Topić (2005) and Flora Croatica Database (Nikolić, 2013a). The new taxa for the Hvar archipelago are marked with the symbol (*) before the name of taxon in the floristic list.

Herbarium specimens are deposited in ZAGR herbarium and are accessible through the Virtual Herbarium ZAGR according to Bogdanović (2013).

Results and discussion

During the floristic research of Zečevo islet a total of 216 vascular plant species and subspecies were found (Tab. 1), out of which 12 species are new records for Hvar Archipelago. According to taxonomical analysis, the dominant group of *Magnolianae* (168 species and subspecies) are almost four times more represented than *Lilianae* (43 species and subspecies). Researched flora includes 65 families, 161 genera, 208 species and 8 subspecies (Tab. 2). The most dominant families are *Fabaceae* with 28 species and subspecies (13%), followed by *Asteraceae* s.l. with 20 species (9.2%), *Poaceae* with 18 species and subspecies (8.3%) and *Lamiaceae* with 11 species (5.1%). Other families are represented with a smaller number of taxa. On the researched area five endangered species were found: *Ophrys bertolonii* Moretti, *O. sphegodes* Mill. and *Salsola kali* L. (VU), *Glaucium flavum* Crantz and *Carex extensa* Gooden. (EN). If the islet of Zečevo compared with other south Adriatic islands and islets, there is a high similarity with respect to plant diversity (Tab. 3). The island of Hvar covers an area of 299.7 km² (Rubić, 1952) and a total of 1095 species and subspecies were recorded (Trinajstić, 1993). Compared to islet of Zečevo, island of Hvar has 3000 times bigger surface, but it has five times more plant taxa. Comparative

Table 1. List of vascular flora for Zečevo islet

Taxon name	Life form	Floral element	Status	<i>Silene vulgaris</i> (Moench) Garcke	H	sem
				<i>Stellaria media</i> (L.) Vill.	T	wp
Chenopodiaceae				<i>Chenopodium murale</i> L.	T	ma
				<i>Salsola kali</i> L.	T	wp
Cichoriaceae				<i>Crepis foetida</i> L.	T	sem
				<i>Crepis neglecta</i> L.	T	eum
				<i>Crepis rubra</i> L.	T	cm
				<i>Hedypnois cretica</i> (L.) Dum.Cours.	T	cm
				<i>Hyoseris radiata</i> L.	H	cm
				<i>Hyoseris scabra</i> L.	T	cm
				<i>Leontodon taraxacoides</i> (Vill.) Mérat	H	mp
				<i>Reichardia picroides</i> (L.) Roth	H	cm
				<i>Sonchus asper</i> (L.) Hill	T	ea
				<i>Sonchus oleraceus</i> L.	T	wp
				* <i>Taraxacum laevigatum</i> auct. croat.	H	se
				<i>Tragopogon porrifolius</i> L.	H	cm
				<i>Urospermum picroides</i> (L.) Scop. ex F.W.Schmidt	T	cm
Cistaceae				<i>Fumana thymifolia</i> (L.) Spach ex Webb	Ch	cm
				Convolvulaceae	H	cm
				<i>Convolvulus cantabrica</i> L.	P	ea
				Cornaceae		
				* <i>Cornus mas</i> L.		
				Corylaceae		
				<i>Ostrya carpinifolia</i> Scop.	P	se
				Crassulaceae		
				<i>Sedum acre</i> L.	Ch	eu
				<i>Sedum rubens</i> L.	T	se
				Cuscutaceae		
				<i>Cuscuta australis</i> R. Br.	T	se
				Dioscoreaceae	G	cm
				<i>Tamus communis</i> L.		
				Ericaceae		
				<i>Arbutus unedo</i> L.	P	cm
				<i>Erica arborea</i> L.	P	cm
				Euphorbiaceae		
				<i>Euphorbia exigua</i> L.	T	cm
				<i>Euphorbia fragifera</i> Jan	Ch	ila
				<i>Euphorbia peplodes</i> Gouan	T	cm
				<i>Euphorbia peplus</i> L.	T	se
				<i>Euphorbia spinosa</i> L.	Ch	cm
				<i>Mercurialis annua</i> L.	T	wp
				Fabaceae		
				<i>Anthyllis vulneraria</i> L. ssp. <i>praeproperta</i> (A.Kern.) Bornm.	H	cm
				<i>Astragalus muelleri</i> Steud. et Hochst.	H	ila
				<i>Ceratonia siliqua</i> L.	P	cm
				<i>Colutea arborescens</i> L.	P	cm
				<i>Dorycnium hirsutum</i> (L.) Ser.	Ch	cm
				<i>Hippocratea ciliata</i> Willd.	T	cm
				<i>Lathyrus cicera</i> L.	T	cm
				<i>Lathyrus setifolius</i> L.	T	cm
				<i>Lathyrus sphaericus</i> Retz.	T	cm
				<i>Lotus corniculatus</i> L.	H	wp
				<i>Lotus cytisoides</i> L.	Ch	cm
				<i>Lotus edulis</i> L.	T	cm
				<i>Lotus ornithopodioides</i> L.	T	cm
				<i>Medicago littoralis</i> Rohde ex Loisel.	T	cm
				<i>Medicago lupulina</i> L.	T	ea
				<i>Medicago minima</i> (L.) Bartal.	T	wp
				<i>Medicago polymorpha</i> L.	T	eum
				<i>Melilotus sulcatus</i> Desf.	T	cm
				<i>Ononis ornithopodioides</i> L.	T	cm
				<i>Ononis reclinata</i> L.	T	cm
				<i>Scorpiurus muricatus</i> L.	T	cm
				* <i>Tetragonolobus maritimus</i> (L.) Roth	H	mp
				<i>Trifolium campestre</i> Schreber	T	wp
				<i>Trifolium scabrum</i> L.	T	cm

<i>Vicia cracca</i> L.	H	ea	<i>Ranunculaceae</i>	G	cm
<i>Vicia lutea</i> L.	T	sem	<i>Anemone hortensis</i> L.	T	cm
<i>Vicia sativa</i> L.	T	ea	<i>Nigella damascena</i> L.	G	eu
<i>Vicia tetrasperma</i> (L.) Schreber	T	se	* <i>Ranunculus ficaria</i> L.		
Fagaceae			<i>Resedaceae</i>		
<i>Quercus ilex</i> L.	P	cm	<i>Reseda lutea</i> L.	H	eu
Fumariaceae			<i>Reseda phyteuma</i> L.	T	cm
<i>Fumaria capreolata</i> L.	T	cm	<i>Rosaceae</i>		
<i>Fumaria officinalis</i> L.	T	ea	<i>Crataegus monogyna</i> Jacq.	P	ea
<i>Fumaria parviflora</i> Lam.	T	sem	<i>Prunus mahaleb</i> L.	P	sep
Gentianaceae			<i>Sanguisorba minor</i> Scop. ssp. <i>muricata</i>	H	sem
<i>Blackstonia perfoliata</i> (L.) Huds.	T	ma	Briq.		
<i>Centaurium erythraea</i> Rafn	H	wp	<i>Rubiaceae</i>		
* <i>Centaurium tenuiflorum</i> (Hoffmanns. et Link) Fritsch	T	ea	<i>Galium aparine</i> L.	T	ea
Geraniaceae			<i>Galium corrudifolium</i> Vill.	H	mp
<i>Erodium cicutarium</i> (L.) L'Hér.	T	wp	<i>Galium lucidum</i> All.	H	sem
<i>Erodium malacoides</i> (L.) L'Hér.	T	cm	<i>Rubia peregrina</i> L.	P	cm
<i>Geranium columbinum</i> L.	T	ea	<i>Sherardia arvensis</i> L.	T	wp
<i>Geranium dissectum</i> L.	T	wp	<i>Valantia muralis</i> L.	T	cm
<i>Geranium molle</i> L.	T	wp	Santalaceae		
<i>Geranium rotundifolium</i> L.	T	wp	<i>Osyris alba</i> L.	G	cm
<i>Geranium robertianum</i> L.	T	wp	Saxifragaceae		
Clusiaceae			<i>Saxifraga tridactylites</i> L.	T	cm
<i>Hypericum perforatum</i> L.	H	ea	Scrophulariaceae		
Lamiaceae			<i>Bellardia trixago</i> (L.) All.	T	cm
* <i>Ajuga genevensis</i> L.	H	ea	<i>Kickxia elatine</i> (L.) Dumort.	T	cm
<i>Lamium amplexicaule</i> L.	T	wp	<i>Verbascum phoeniceum</i> L.	H	sep
<i>Marrubium incanum</i> Desr.	H	cm	<i>Veronica cymbalaria</i> Bodard	T	cm
<i>Micromeria graeca</i> (L.) Rchb.	Ch	cm	Solanaceae		
<i>Micromeria juliana</i> (L.) Benth. ex Rchb.	Ch	cm	<i>Solanum nigrum</i> L.	T	wp
<i>Prasium majus</i> L.	Ch	cm	Thymelaeaceae		
<i>Rosmarinus officinalis</i> L.	P	cm	* <i>Thymelaea hirsuta</i> (L.) Endl.	P	cm
<i>Sideritis romana</i> L.	T	cm	Valerianaceae		
<i>Teucrium polium</i> L.	Ch	mp	<i>Valerianella echinata</i> (L.) DC.	T	cm
<i>Thymus longicaulis</i> C.Presl	Ch	ila	Verbenaceae		
<i>Thymus serpyllum</i> L.	Ch	eu	<i>Vitex agnus-castus</i> L.	P	cm
Linaceae			Violaceae		
<i>Linum strictum</i> L. ssp. <i>corymbulosum</i>	T	cm	<i>Viola arvensis</i> Murray	T	wp
(Rchb.) Riony			Lilianeae		
<i>Linum trigynum</i> L.	T	cm	* <i>Narcissus tazetta</i> L.	G	cm
Loranthaceae			* <i>Narcissus serotinus</i> L.	G	cm
* <i>Loranthus europaeus</i> Jacq.	P	ea	Araceae		
Malvaceae			<i>Arisarum vulgare</i> O.Targ.Tozz.	G	cm
<i>Althaea hirsuta</i> L.	T	cm	<i>Arum italicum</i> Mill.	G	cm
<i>Lavatera arborea</i> L.	H	cm	Asparagaceae		
<i>Malva sylvestris</i> L.	H	ea	<i>Asparagus acutifolius</i> L.	G	cm
Moraceae			Cyperaceae		
<i>Ficus carica</i> L.	P	eum	<i>Carex distachya</i> Desf.	H	cm
Myrtaceae			<i>Carex extensa</i> Gooden.	H	ma
<i>Myrtus communis</i> L.	P	cm	<i>Carex hallerana</i> Asso	H	cm
Oleaceae			Iridaceae		
<i>Fraxinus ornus</i> L.	P	eum	<i>Iris germanica</i> L.	H	wp
<i>Olea europaea</i> L.	P	cm	<i>Romulea bulbocodium</i> (L.) Sebast. et	G	cm
<i>Phillyrea media</i> L.	P	cm	Mauri		
Orobanchaceae			Liliaceae		
<i>Orobanche minor</i> Sm.	T	sem	<i>Allium subhirsutum</i> L.	G	cm
Papaveraceae			<i>Asphodelus aestivus</i> Brot.	G	cm
<i>Glaucium flavum</i> Crantz	H	cm	<i>Bellevalia dubia</i> (Guss.) Rchb.	G	cm
Plantaginaceae			<i>Muscaris comosum</i> (L.) Mill.	G	sem
<i>Plantago afra</i> L.	T	cm	<i>Muscaris neglectum</i> Guss. ex Ten.	G	cm
<i>Plantago holosteum</i> Scop.	H	cm	<i>Ornithogalum collinum</i> Guss.	G	cm
Plumbaginaceae			<i>Ornithogalum comosum</i> L.	G	mp
<i>Limonium cancellatum</i> (Bernh. ex Bertol.)	H	ila	<i>Ornithogalum umbellatum</i> L.	G	sem
Kuntze		e	<i>Scilla autumnalis</i> L.	G	mp
<i>Limonium narbonense</i> Mill.	H	cm	<i>Smilax aspera</i> L.	G	cm
Portulacaceae			Orchidaceae		
<i>Portulaca oleracea</i> L.	T	wp	<i>Limodorum abortivum</i> (L.) Sw.	G	cm
Primulaceae			<i>Ophrys bertolonii</i> Moretti	G	sem
<i>Anagallis arvensis</i> L.	T	wp	<i>Ophrys liburnica</i> Devillers et Devillers-	G	ila
<i>Anagallis coerulea</i> Schreb.	T	cm	Tersch.		VU e

<i>Ophrys sphegodes</i> Mill.	G	cm	VU
<i>Serapias parviflora</i> Parl.	G	cm	
Poaceae			
<i>Anthoxanthum odoratum</i> L.	H	ea	
<i>Avena barbata</i> Pott ex Link	T	ma	
<i>Brachypodium retusum</i> (Pers.) P.Beauv.	H	cm	
<i>Briza maxima</i> L.	T	wp	
<i>Bromus hordeaceus</i> L.	T	wp	
<i>Bromus madritensis</i> L.	T	ma	
* <i>Catapodium pauciflorum</i> (Merino)	T	cm	
Brullo, Giusto, Miniss. & Spamp.			
<i>Cynodon dactylon</i> (L.) Pers.	G	wp	
<i>Dactylis glomerata</i> L.	H	wp	
<i>Desmazeria rigida</i> (L.) Tutin	T	ma	
<i>Heteropogon contortus</i> (L.) P.Beauv. ex Roem. et Schult.	H	cm	NT
<i>Hordeum murinum</i> L. ssp. <i>leporinum</i> (Link) Arcang.	T	cm	
<i>Koeleria splendens</i> C.Presl	H	sem	
<i>Lagurus ovatus</i> L.	T	cm	
<i>Lolium rigidum</i> Gaudin	T	sem	
<i>Lophochloa cristata</i> (L.) Hyl.	T	ea	
<i>Melica ciliata</i> L.	H	cm	
<i>Poa bulbosa</i> L.	H	wp	

Table 2. Taxonomical analysis of the flora of Zečevo islet

Taxon	Monoliophyta	Gymnospermae	Angiospermae		Σ
			Magnolianae	Lilianae	
Family	1	3	54	7	65
Genus	1	3	123	34	161
Species	1	3	162	42	208
Subspecies	0	1	6	1	8
Species and subspecies (%)	0.5	1.9	77.8	19.9	100.0

Table 3. Number of taxa and area size of the islet of Zečevo (Hvar) and some middle and south Adriatic islands and islets (1, 3, 4, 6: Regula-Bevilacqua and Ilijanić, 1984; 2: Trinajstić, 1993; Nikolić, 2013a; 5: Bogdanović et al., 2004; 7: Milović, 2004; 8: Vladović et al. 2001; 9: present study; 10: Jasprica et al., 2006; 11: Pavletić, 1983)

Island	Area (km ²)	Number of species and subspecies
1. Brač	394.57	750
2. Hvar	299.70	1224 (with new data)
3. Korčula	276.00	850
4. Vis	90.30	800
5. Biševo	5.80	421
6. Svetac	5.10	344
7. Obonjan (Šibenik)	0.70	230
8. Vela Kluda (Trogir)	0.12	137
9. Zečevo (Hvar)	0.11	216
10. Sv. Andrija (Dubrovnik)	0.05	160
11. Brusnik	0.04	41

analysis between Zečevo and Hvar archipelago (Trinajstić, 1993; Nikolić, 2013a) results in identification of 12 new species (5.6% of the total flora registered on Zečevo islet) that have not been found on the island of Hvar before.

Analysis of life forms shows a high ratio of therophytes (45.4%) and hemicryptophytes (21.3%) (Fig. 2). The therophytes dominance indicates influence of typical Mediterranean climate on the islet. The high percentage of geophytes (12.5%) indicates floristically rich spring aspect of the flora of Zečevo islet. It is important to note that among the geophytes there are some vulnerable (VU) plant species as: *Ophrys bertolonii* Moretti and *Ophrys sphegodes* Mill. and near threatened (NT) like *Narcissus tazetta* L. The spectrum of life forms of Zečevo islet coincides with the spectra of some similar size area islands and islets of central and southern Adriatic as well as with the spectrum of the Mediterranean in general (Tab. 4). Compared with spectrum of the Mediterranean in general (Horvat, 1949), there is some larger proportion of therophytes and smaller proportion of hemicryptophytes in the spectrum of Zečevo. A slightly larger proportion of therophytes is probably because the islet of Zečevo has a lot of open grassland habitats.

Phytogeographical analysis (Fig. 3) showed prevalent of Mediterranean floral element (138 species and subspecies, 63.4%) of which the most of circum Mediterranean plants (111 species and subspecies, 51.4%) followed by widespread plants represented by 27 species (12.5%) and plants of Euro-Asian floral element with 21 species and subspecies (9.7%). Other floral elements were represented by smaller number. Significant presence of Mediterranean floral element indicates that Zečevo islet belongs to the corresponding Mediterranean region (cf. Horvatić, 1963)




Figure 2. Analysis of life forms on Zečevo islet (T - Therophyta, H - Hemicryptophyta, P - Phanerophyta, G - Geophyta and Ch - Chamaephyta)




Figure 3. Analysis of floral elements in the flora of Zečevo islet

Table 4. Comparative survey of life forms of Zečevo islet and similar size area middle and south Adriatic islands, islets and the Mediterranean (%)

Area	T (%)	H (%)	P (%)	Ch (%)	G (%)	Literature source
Obonjan	47	19.1	11.7	11.7	10.4	Milović, 2004
Vela Kluda	45.3	21.2	13.1	8.8	11.6	Vladović et al., 2001
Zečevo	45.4	21.3	13	7.9	12.5	present study
Sv. Andrija	36.4	25.3	18.2	13	6.5	Jasprica et al., 2006
Mediterranean	42	29	12	6	11	Horvat, 1949

Table 5. Comparative survey of the prevalent floral elements of Zečevo islet and some other similar size areas of Adriatic islets (%)

Area	Mediterranean	South European	Euro-Asian	Widespread plants	Alien	Literature source
Obonjan	53	20.4	3.9	16.1	3	Milović, 2004
Vela Kluda	63.5	17.5	3	13.8	1.5	Vladović et al., 2001
Zečevo	63.4	7.4	9.7	12.5	1.9	present study
Sv. Andrija	46.2	15.8	6.3	18.4	11.4	Jasprica et al., 2006

**Figure 4.** *Ophrys liburnica* Devillers et Devillers-Tersch (photo: M. Skelin)

as expected. Share of Mediterranean floral element of Zečevo corresponds to the islets of similar size in central and southern Adriatic (Tab. 5). In contrast wider variations in the share of South European and Euro-Asian floral element were established. Smaller proportion of alien taxa can be explained by low anthropogenic influence on the islet, although a portion of ruderal and nitrophilous plants (as widespread plants) are high because of the presence of seagulls and rabbits on the islet. Within the group of the Illyrian-Adriatic endemic plants five taxa were recorded: *Vincetoxicum hirundinaria* Medik. ssp. *adriaticum*

**Figure 5.** *Narcissus serotinus* L. (photo: M. Skelin)

(Beck) Markgr., *Aurinia sinuata* (L.) Griseb., *Limonium cancellatum* (Bernh. ex Bertol.) Kuntze., *Astragalus muelleri* Steud. et Hochst. and *Ophrys liburnica* Devillers et Devillers-Tersch.

The flora of Zečevo islet is particularly valuable because of the presence of three rare plant species (*Ophrys liburnica* Devillers et Devillers-Tersch, *Scaligeria cretica* (Mill.) Boiss. and *Narcissus serotinus* L.) of the Croatian flora. New localities of these species are important in chorological point of view as the distribution area for these species is expanded outside known localities. Few individuals of stenoendemic orchid *Ophrys liburnica* Devillers et Devillers-Tersch (Fig. 4) were found in the *Pinus halepensis* wood. This species was known from Istria, Quarner Archipelago (Devillers and Devillers-Terschuren, 2004) from Korčula, Pelješac peninsula, Hvar and some costal parts of central Dalmatia area of Šibenik and Split (Delforge 2006). *Scaligeria cretica* (Mill.) Boiss. as a representative of *Apiaceae* has white flowers grouped in umbels and can often be confused with other similar taxa of this family. So far, this taxon has been recorded for some central and southern Adriatic islands and islets: Svetac, Vis, Brač, Hvar, Šćedro and in Molunat (Konavle area) (Visiani, 1852; Horvatić,

1970; Hodak, 1971; Pavletić, 1978; Trinajstić, 1993). The finding of *S. cretica* on Zečevo islet is expected due to relative vicinity of the island of Hvar, where this species is common (Trinajstić, 1993). As an autumn flowering species of daffodil, *Narcissus serotinus* L. (Fig. 5) is often omitted from floristic lists of some areas of Croatian territory. According to the latest overview of its distribution in Croatia *N. serotinus* is present on Ugljan, Promina, Knin area (Devetak, 2000), Lastovo, Pelješac peninsula and Prevlaka in Konavle region (Nikolić, 2013a).

Conclusion

In vegetation period from 2006 to 2010 flora mapping of Zečevo islet was performed and the total number of 216 vascular plant species and subspecies were recorded, while 12 species were recorded as a new for the Hvar archipelago. Prevalent of Mediterranean floral element and therophytes dominance indicates influence of typical Mediterranean climate on the islet. Compared with other middle and south Adriatic islands, the island of Hvar has the highest number of vascular plant species. Given that, on the islet of Zečevo was found several rare and endemic plant species.

References

- Bogdanović S., Mitić B. (2003). The flora of the volcanic island of Brusnik (central Dalmatia, Croatia). *Acta Bot. Croat.* 62(2): 103-113
- Bogdanović S., Dobrović I., Ostojoć A., Boršić I., Modrić Ž., Vojnić Rogić I. (2004). A contribution to the vascular flora of the island of Biševo (central Adriatic, Croatia). *Acta Bot Croat* 63(2): 125-134
- Bogdanović S., ed. (2013). Virtual Herbarium ZAGR. University of Zagreb, Faculty of Agriculture. <http://herbarium.agr.hr> (accessed: 02 October 2013)
- Boršić I., Milović M., Dujmović I., Bogdanović S., Cigić P., Rešetnik I., Nikolić T., Mitić B. (2008). Preliminary check-list of invasive alien plant species in Croatia. *Nat Croat* 17(2): 55-71
- Delforge P. (2006). Contribution à la connaissance des Orchidées de Croatie. Résultats de cinq années de prospections. *Natural belges* 87(Orchid. 19):141-200
- Devetak Z. (2000). A new locality of the species *Narcissus serotinus* L. - autumn daffodil in Croatia. *Nat Croat* 9(2):157-162
- Devillers P., Devillers-Terschuren J. (2004). The *Ophrys sphegodes* complex in the Adriatic: spatial and temporal diversity. *Natural belges* 85 (Orchid. 17):129-148
- Duplančić Leder T., Ujević T., Čala M. (2004). Duljine obalne crte i površine otoka na hrvatskom dijelu Jadranskog mora određene s topografskih karata mjerila 1:25 000. *Geoadria* 9(1): 5-32
- Hodak N. (1971). Neka nova nalazišta vrste *Scaligeria cretica* (Miller) Boiss. na području hrvatskog primorja. *Acta Bot Croat* 30:119-121
- Horvat I. (1949). Nauka o biljnim zajednicama. Nakladni zavod Hrvatske, Zagreb
- Horvatić S. (1963). Vegetacijska karta otoka Paga s općim pregledom vegetacijskih jedinica Hrvatskog primorja. *Priroda Jugosl Akad* 33. *Acta Biolog* 4:1-187
- Horvatić S. (1970). *Scaligeria cretica* (Miller) Boiss. u bilnjem pokrovu hrvatskog primorja. *Acta Bot Croat* 29:175-182
- Jasprica N., Kovačić S. (1997). Vascular flora of the central part of Pelješac peninsula. *Nat Croat* 6(4): 381-407
- Jasprica N., Kovačić S., Ruščić M. (2006). Flora and vegetation of Sveti Andrija island, southern Croatia. *Nat Croat* 15(1-2):27-42
- Gabelica I., Piasevoli G., Jurić M., Mekinić S., Kažimir Z., Pešić N., Perković A. (2011). Zaštićene prirodne vrijednosti Splitsko-dalmatinske županije. Javna ustanova za upravljanje zaštićenim prirodnim vrijednostima na području Splitsko-dalmatinske županije, Split
- Kovačić S., Jasprica N., Lasić T. (2000). A contribution to the vascular flora of the Pelješac peninsula (southern Croatia). *Acta Bot Croat* 59(2): 411-419
- Mihovilović M. A., ed. (1995). *Otok Hvar, Otoci*, Matica Hrvatska, Zagreb
- Milović M. (2004). Flora of the island of Obonjan (Šibenik archipelago, Croatia). *Nat Croat* 13(3):213-230
- Nikolić T., Topić J. (2005). Crvena knjiga vaskularne flore Republike Hrvatske. Kategorije EX, RE, CR, EN i VU. Ministarstvo kulture, Državni zavod za zaštitu prirode, Zagreb
- Nikolić T. (2006). Priručnik za inventarizaciju i praćenje stanja. Državni zavod za zaštitu prirode, Zagreb
- Nikolić T., ed. (2013a). Flora Croatica Database. <http://hirc.botanic.hr/fcd> (accessed: 02 October 2013)
- Nikolić T. (2013b). Sistematska botanika: Raznolikost i evolucija biljnog svijeta. ALFA d.d., Zagreb
- Pavletić Zi. (1978). Pregled i analiza flore Palagruških otoka. *Biosistematika* 4(1):39-47
- Pavletić Zi. (1983). Pregled flore i vegetacije nekih manjih srednjedalmatinskih otoka i otočića. In: Pavletić Z., Matković P., Grubišić S. (eds.): *Zbornik Roberta Visianija Šibenčanina (Povremena izdanja Muzeja grada Šibenika, Svezak 10)*. Muzej grada Šibenika, Šibenik, 315-329
- Pignatti S. (1982). *Flora d'Italia I-III*. Edagricole, Bologna
- Raunkiaer C. (1934). The Life Forms of Plants and Statistical Plant Geography, being the collected papers of C. Raunkiaer. Reprinted 1978 (ed. by Frank N. Egerton), Oxford University Press, Oxford
- Regula-Bevilacqua Lj., Ilijanić Lj. (1984). Analyse der Flora der Insel Mljet. *Acta Bot Croat* 43:119-142
- Rubić I. (1952). Naši otoci na Jadranu. Izdanje Odbora za proslavu desetgodišnjice Mornarice, Split
- Trinajstić I. (1993). Vaskularna flora otoka Hvara. *Acta Bot Croat* 52:113-143
- Trinajstić I. (1998). Supplements to the flora of island of Korčula (Croatia). *Acta Bot Croat* 57:95-98
- Visiani R. (1852). *Flora Dalmatica*. Vol. 3. Apud Friedericum Hofmeister, Lipsiae
- Vladović D., Jukić N. (1997). Prilog flori otoka Korčule. *Acta Bot Croat* 55/56:139-141
- Vladović D., Šunjara V., Pavlov V., Bačić T. (2001). Vascular flora of the island of Vela Kluda. *Nat Croat* 10(1):19-31
- Vladović D., Mitić B., Martinović P. (2002). A contribution to the flora of the island of Mali Drvenik. *Nat Croat* 11(2):237-242