

NEW RECORD OF MIDDLE JURASSIC RADIOLARIANS AND EVIDENCE OF NEOTETHYAN DYNAMICS DOCUMENTED IN A MÉLANGE FROM THE CENTRAL DINARIDIC OPHIOLITE BELT (CDOB, NE BOSNIA AND HERZEGOVINA)

Branimir Šegvić*, **Duje Kukoč****, **Ivan Dragičević*****, **Alan Vranjković*****, **Vlatko Brčić°**, **Špela Goričan****, **Elvir Babajić°°** and **Hazim Hrvatović°°°**

* *Section des Sciences de la Terre et de l'Environnement, Université de Genève, Switzerland.*

** *Paleontološki Inštitut Ivana Rakovca ZRC SAZU, Ljubljana, Slovenia.*

*** *Faculty of Mining, Geology and Petroleum Engineering, University of Zagreb, Croatia.*

° *Croatian Geological Survey, Zagreb, Croatia.*

°° *Faculty of Mining, Geology and Civil Engineering, University of Tuzla, Bosnia and Herzegovina.*

°°° *Geological Survey of Federation of Bosnia and Herzegovina, Sarajevo-Ilidža, Bosnia and Herzegovina.*

✉ *Corresponding author email: branimir.segvic@unige.ch*

Keywords: *Jurassic radiolarians, ophiolitic mélangé, Dinarides, CDOB, Neotethyan dynamics. Bosnia and Herzegovina.*

ABSTRACT

Within the ophiolitic mélangé of the Central Dinaridic Ophiolitic Belt (CDOB) that stretches throughout the Balkans region in SE Europe, a latest Bajocian-early Bathonian radiolarian assemblage was obtained from chert-rich shaly to silty matrix. The sampling locality in northern Bosnia and Herzegovina is characterized by a highly-diversified ophiolitic suite, consisting of basic and ultrabasic rocks of different geotectonic provenances. This makes the radiolarian dating a convenient complementary tool for studying the geodynamic history of CDOB within a broader regional context. The host sediments and the nature of their associated crystalline rocks suggest that radiolarian deposition occurred relatively close to the Adria shelf margins, predating or being contemporaneous to the rapid transitions in the Dinaridic Neotethys geotectonic setting, changing from active ridge magmatism to an intraoceanic subduction environment and island-arc volcanism. The minimum age of ophiolite mélangé formation is defined by the mineral equilibration ages in metamorphic sole (161 ± 4 Ma), with the obduction tectonics that must have lasted at least until the Oxfordian time (i.e. termination of MOR activity in the Dinarides). This age correlates well with the ages of sediments reported elsewhere in the mélangé of the Dinaride-Hellenide orogenic system.

INTRODUCTION


The Jurassic ophiolites found in narrow zones of the Dinarides are part of the Alpine-Himalayan-Tethyan orogenic belt and are characterised by two belts of ophiolites: (1) the Central Dinaridic Ophiolite Belt (CDOB) in the west, and (2) the Vardar Zone (VZ) in the east (Fig. 1). These can be continuously traced to the south into Albania, Macedonia and Greece. Both belts are primarily composed of ophiolitic mélangé with ophiolitic blocks and various genetically-related sedimentary rocks (e.g., Dimitrijević et al., 2003). Previously believed that such defined ophiolite domains stood for two discrete Neotethyan oceanic branches (e.g., Spray et al., 1984; Smith and Spray, 1984; Lugović et al., 1991; Robertson et al., 2013), the approaches nowadays advocate the origin of Dinaridic ophiolites from the very same palaeogeographic oceanic branch of Neotethys with a mechanism of westwards sheet thrusting that conditioned their current, klippe-like, structural position within the Dinaride-Hellenide nappe stack (e.g., Çollaku et al., 1992; Pamić et al., 1998; Gawlick et al., 2008; Schmid et al., 2008; Ustaszewski et al., 2010).

Starting in the Middle or late Early Jurassic, the oceanic realm partly preserved in CDOB and VZ experienced an intraoceanic subduction, resulted from the latest Mesozoic to Cenozoic SW-vergent tectonics, following the collision between Adria and Europe-derived units. The initiation of convergence gave rise to incipient intraoceanic subduction and metamorphism at the ophiolite base (e.g., Operta et al., 2003; Karamata, 2006; Šegvić, 2010). Impelled by the for-

mer, a system of intraoceanic trenches formed, being progressively filled by gravitational emplacements of chaotic material from the upper plate (radiolarite-ophiolite slide blocks with clayey to sandy matrix) and clastic fragments emanated from the footwall. Due to the constant shortening, tectonically overprinted allochthonous material is reworked into the accretionary prisms, thus defining a widely accepted mechanism of mélangé formation in the Dinaride-Hellenide orogeny (e.g., Gawlick et al., 2008; 2009; Bortolotti et al., 1996; 2004). Ultimately, the deposition of ophiolitic mélangé evolved with the onward obduction processes of the oceanic lithosphere, with a composite thrust sheets system being created, whereby the mélangé is positioned in the footwall of the main ultramafic mass of CDOB (e.g., Pamić et al., 2002; general SW-NE profile in Fig. 2, Dragičević, 1987). Still, the exact way of the Middle Jurassic mélangé formation in the Dinarides is poorly known, especially in regard to the provenance of its components and the geotectonic constraints inferred thereof (e.g., Gawlick et al., 2009). Detailed case studies focusing on the age and the composition of the CDOB mélangé are thus necessary to complement existing efforts done elsewhere in the Dinarides-Hellenides, shaping a larger picture on the palaeogeography of mélangé formation. Biostratigraphic data exist from the ophiolitic mélangé in Croatia (Halamić et al. 1999), Serbia (Gawlick et al. 2009; Chiari et al. 2011), Albania (Mirdita Zone, e.g. Kellici et al. 1994; Marcucci et al. 1994; Chiari et al. 2004; Gawlick et al. 2008), and Greece (Danelian and Robertson 2001; Bortolotti et al. 2008; Nirta et al. 2010;



Fig. 1 - Schematic geotectonic map of SE Europe, showing the stretching of the Alpine ophiolite complexes throughout the Balkans. Location of sampling indicated by rectangle. Map modified after Robertson (2002).



Chiari et al. 2012). All obtained ages are coherent, yielding a Middle Jurassic age of mélangé. So far, Middle Jurassic radiolarians in the Bosnian part of CDOB were described only from non-ophiolitic sequences (Vishnevskaya et al., 2009). In this paper, new evidence is presented on the age of Jurassic radiolarian chert recovered from the mélangé of the biggest ophiolite complex in the central CDOB, the Krivaja-Konjuh Ophiolite Complex (KKOC, NE Bosnia and Herzegovina), in which diverse oceanic rock suites testify to the different stages of ocean evolution (Pamić et al., 1977; Lugović et al., 1991; 2006; Babajić, 2009; Šegvić, 2010). Obtained radiolarian ages along with the mélangé composition and isotopic ages of its crystalline components are discussed in terms of their significance for the Jurassic Neotethyan evolution in its Dinaridic segment.

GEOLOGICAL SETTING

The Krivaja-Konjuh Ophiolite Complex (of about 650 km² in northern Bosnia and Herzegovina (Fig. 1, Pamić and Hrvatović, 2000). The two KKOC structural subunits - Krivaja, the western subunit consisting of amalgamated peridotite blocks, and Konjuh, the eastern coherent peridotite synform - are thrust onto the ophiolitic mélangé and overstepped by the Cretaceous basinal Pogari Formation (Pamić,

Fig. 2 - Composite stratigraphic section of the Papratnica section. Sampling position indicated within the sequence. Constructed after basic geological map of Yugoslavia, sheet Vareš (Pamić, 1970). Time scale after Gradstein et al. 2012.

1970; Hrvatović, 2006). Ophiolite mélangé forms the largest part of the KKOC (Fig. 3). It presents a chaotic tectono-sedimentary mixture made up of detached 'pocket' to 'mountain'-size blocks and boulders of oceanic crust rocks along with Palaeozoic to Jurassic carbonates and clastics (Tari, 2002). More precisely, the lithology of the mélangé comprises a sheared pelitic to silty matrix in which greywacke fragments predominate over basalt and tuff, diabase, gabbro, serpentized peridotite with metamorphic rocks, shale, radiolarite, and exotic blocks of recrystallized pseudo-oolitic and nodular Middle to Late Triassic to Late Jurassic (Tithonian) limestones and dolomite limestones (Pamić, 1968; 1982, Fig. 2). Poorly preserved microfossils of Jurassic foraminifera (*Vidalina martana*, *Glomospira* sp., *Cristellaria* sp.) are reported extremely rarely in micrites interlayered with shale and greywacke (Pamić and Hrvatović, 2000). This roughly defines the Jurassic age of the KKOC (Pamić and Hrvatović, 2000). The K-Ar isotopic ages of 174-157 Ma from amphiboles separated from the metamorphic base favour the latter (Lanphere et al., 1975; Majer et al., 1979). Breccias and conglomerates of the overstepping Cretaceous succession contain ophiolitic fragments and, thus constrain the youngest possible age of the Bosnian CDOB mélangé (Fig. 3, Jovanović et al., 1978, Pamić and Hrvatović, 2000; Hrvatović, 2006, and references therein).

The ophiolite mélangé was studied and sampled in the vicinity of the village Papratnica and the homonymous creek located east of Žepče in northern Bosnia and Herzegovina (locality no. 1 in Fig. 3, GPS coordinates N 44°25'1.95" E 17°59'0.39"). The sampling area presents the NW peak segment of the KKOC, which is detached from the main ophiolite mass. The Papratnica mélangé deposits were selected for sampling because they comprise carbonate blocks and pillow lavas, hence being roughly correlated to the olistostrome part of the CDOB ophiolitic mélangé (Pamić, 1968). In stratigraphical sense, the base of Papratnica section commences with tens of metres thick, highly altered bodies of pillow lavas, having no direct contact with radiolarian cherts (Fig. 2). Apparently, the former are continuously overlain by the greenish-black to reddish-black sheared shales intercalated with mudstones, siltites, and sandstones. The section ends with a several metres wide carbonate block (Fig. 4b). The nearly 100 m thick Papratnica mélangé succession holds numerous clast horizons containing radiolarian chert, one of which was logged and sampled at the second metre of the section. This outcrop sampled for radiolarians is approximately 4 m thick and consists of layered dark grey fine-grained shales with embedded chert blocks (Fig. 4a). The proportion of pelitic interlayers is relatively high, accounting for approximately 30% of the




Fig. 3 - Detailed geological map of the sampling area within the western portion of the Krivaja-Konjuh ophiolite complex (KKOC). 1- sampling location, this work; 2- sampling location of the non-ophiolitic chert succession described in Vishnevskaya et al. (2009). Map modified after Jovanović et al. (1978), Pamić (1970), and Dragičević (1987).




Fig. 4 - a) layered dark grey fine-grained shales with embedded chert blocks; b) the large limestone olistolith that tops the Papratnica section.

sampled section. The succession is pervasively sheared and, as a result of shearing, the siliceous-shaly beds are disrupted, boudinaged, and some of them brecciated. Due to the complex and chaotic nature of *mélange*, it was not possible to log a complete Papratnica section, which means that the position of studied radiolarians within the *mélange* column is approximate (Fig. 2).

The black and vitreous subangular to rounded chert clasts vary from 2 to 10 cm in diameter, whereas in some other levels of the Papratnica section they are up to 20 cm in diameter. Thin-section analyses showed that the sampled clasts consist of angular chert fragments (0.5-2 cm) that are imbedded in the cryptocrystalline siliceous matrix. Both the former and the latter contain circular or elliptical areas (ca. 0.05 mm in diameter) filled with microcrystalline quartz. These areas represent complete and nicely preserved radiolarian skeletons. The rock itself is determined as autoclastic radiolarian-chert microbreccia (cf. Kolodny et al., 2005, and references therein). The texture of microbreccia is clast-supported with a medium packing density, whilst the matrix presents a loosely-packed packstone. Such biosiliceous sediments originated from the radiolarian skeleton deposition that characterised large portions of the deep Jurassic Tethys, both in its continental margin and ocean floor settings (e.g., Aiello et al., 2008). No index fossils are reported from the background fine-clastic sedimentary rocks and, therefore, the general Jurassic age of the *mélange* is inferred based on stratigraphically uncharacteristic microfossils found in platy limestones that are interlayering with shales (Pamić et al., 2002). Thin section analyses of the large limestone olistolith topping the Papratnica section (Figs. 2, 4b) revealed a structureless micrite devoid of any recognizable carbonate grains. Based on previous studies of carbonate olistoliths in a wider area (Pamić, 1968; Pamić et al., 1979), we presume that the block is Triassic in age.

RADIOLARIAN DATING

A well-preserved radiolarian assemblage was extracted from a chert sample. The sample was treated with diluted (5%) hydrofluoric acid. The identified radiolarians are listed in Table 1 and illustrated in Plate 1. The assemblage was

dated according to the Unitary Association Zones (UAZ) established by Baumgartner et al. (1995b). The genera designations follow the revision in the Catalogue of Mesozoic radiolarian genera (O'Dogherty et al., 2009).

The radiolarians are characteristic of the uppermost Bajocian-lower Bathonian UAZ 5 of Baumgartner et al. (1995b). This assignment is based on the co-occurrence of *Unuma latusicostatus* (Aita), last appearing in UAZ 5, and *Eucyrtidiellum semifactum* Nagai and Mizutani first appearing in UAZ 5. Another species that, according to Baumgartner et al. (1995b), suggests the upper age limit in UAZ 5 is *Striatojaponocapsa synconexa* O'Dogherty, Goričan and Dumitrica (it was *Tricolocapsa plicarum* ssp. A in Baumgartner et al., 1995a). It is now known that *Striatojaponocapsa synconexa* extends at least to UAZ 6-7 (Prela et al., 2000; O'Dogherty et al., 2006). *Eucyrtidiellum unumaense dentatum* Baumgartner has a stratigraphic range from UAZ 6 to UAZ 7, which is in disagreement with the inferred assignment to UAZ 5. We note, however, that the record of this species in the database of Baumgartner et al., (1995b) is scarce. In addition, the closely related *Eucyrtidiellum unumaense pustulatum* Baumgartner, having a comparatively more complex ornamentation, first occurs in UAZ 5. The range of *Eucyrtidiellum unumaense dentatum* is thus likely to be extended to UAZ 5.

COMPARISON WITH PREVIOUSLY REPORTED RADIOLARIAN AGES IN CORRELATIVE OPHIOLITE COMPLEXES

The latest Bajocian-early Bathonian age of the KKOC is consistent with other ages reported from radiolarian-bearing rocks in ophiolite complexes throughout the Dinarides and Hellenides (Fig. 5). In these complexes, Jurassic radiolarites were found in *mélange* units, as well as in direct stratigraphic contact with basalts. The ophiolitic *mélange* on Mt. Medvednica (NW Croatia), located north westwards of the KKOC, contains radiolarites dated as latest Bajocian - early Bathonian (UAZ 5) and late Bathonian - early Callovian (UAZ 7) (Halamić et al., 1999). Near Sjenica (SW Serbia), Gawlick et al. (2009) reported a Bathonian to early-middle Callovian age from the radiolarian-rich matrix of the

Table 1 - Occurrence of radiolarian taxa in the analysed sample. The second column gives the zone range of species according to Baumgartner et al. (1995b).

Radiolarians	UAZ 95
<i>Archaeodictyomitra rigida</i> Pessagno	-
<i>Archaeodictyomitra suzukii</i> Aita	-
<i>Archaeodictyomitra</i> sp.	-
<i>Canoptum krahsteinense</i> (Suzuki and Gawlick)	-
<i>Dictyomitrella</i> (?) <i>kamoensis</i> Mizutani and Kido	3-7
<i>Eoxitus baloghi</i> Kozur	-
<i>Eucyrtidiellum semifactum</i> Nagai and Mizutani	5-7
<i>Eucyrtidiellum unumaense dentatum</i> Baumgartner	6-7
<i>Helvetocapsa matsukoi</i> (Sashida)	-
<i>Hemicryptocapsa marcucciae</i> (Cortese) (= <i>Williriedellum</i> sp. A of Baumgartner et al. 1995a)	4-8
<i>Hemicryptocapsa yaoui</i> (Kozur)	-
<i>Paronaella broennimanni</i> Pessagno	4-10
<i>Praewilliriedellum</i> sp. aff. <i>P. convexum</i> (Yao)	-
<i>Praewilliriedellum</i> sp. aff. <i>P. himedaruma</i> (Aita)	-
<i>Protunuma lanosus</i> Ožvoldova	-
<i>Quarticella</i> sp.	-
<i>Semihsum amabile</i> (Aita)	4-7
<i>Stichocapsa</i> (?) <i>magnipora</i> Chiari, Marcucci and Prela	-
<i>Stichomitra</i> (?) <i>takanoensis</i> Aita gr.	3-7
<i>Striatojaponocapsa synconexa</i> O'Dogherty, Goričan and Dumitrica (= <i>Tricolocapsa plicarum</i> ssp. A of Baumgartner et al. 1995a)	4-5
<i>Transsum brevicostatum</i> (Ožvoldova) gr.	3-11
<i>Transsum maxwelli</i> (Pessagno) gr.	3-10
<i>Triactoma</i> (?) sp.	-
<i>Unuma</i> sp.	-
<i>Unuma gordus</i> Hull (= <i>Unuma</i> sp. A of Baumgartner et al. 1995a)	4-6
<i>Unuma latusicostatus</i> (Aita)	2-5
<i>Yaocapsa</i> sp. A	-
Age (UAZ 95)	5

mélange. The stratigraphic range of *Hemicryptocapsa tetragona* (Matsuoka), found in sample SCG 84 of Gawlick et al. (2009), is confined to UAZ 5 (Baumgartner et al., 1995b) and, therefore, the corrected age of this sample should be latest Bajocian to early Bathonian (UAZ 5). In the Mirdita Zone in Albania, the ophiolitic mélange ranges in age from the late Bajocian - early Bathonian to the Oxfordian; late Bajocian - early Bathonian radiolarian cherts systematically occur at the base of the chaotic mass-flow deposits (Gawlick et al., 2008). In southern Albania, UAZ 5 radiolarian cherts were found on top of polymictic breccia above basalt (see section Lubonja in Chiari et al., 2004). Radiolarites in direct stratigraphic contact with basalts were also found at several localities and were dated as latest Bajocian - early Bathonian (UAZ 5) to late Bathonian - early Callovian (UAZ 7) (Prela et al., 2000; Chiari et al., 2004). Further south, in Greece, radiolarites of latest Bajocian - early Bathonian (UAZ 5) and middle Bathonian (UAZ 6) ages were documented in the ophiolite complexes of the western ophiolite belt Pindos (UAZ 5 in a block preserving the contact between chert and basalt, Nirta et al., 2010), Othrys (UAZ 6 in chert included in the Agoriani Mélange, Bortolotti et al., 2008) and Koziakas (UAZ 6 in a thick boudinaged chert succession without stratigraphic contact with other formations, Chiari et al., 2012). In the Vourinos Massif, radiolarites of latest Bajocian - early Bathonian

(UAZ 5) to middle Bathonian (UAZ 6) age were found in direct contact with basalts (Chiari et al., 2003). In our short review (Fig. 5), only radiolarian data that restrict the age to a single UAZ are presented. The published ages from several other localities are more broadly defined but also cluster around UAZs 5 to 7. For a complete review of radiolarian ages in ophiolitic units of Albania and Greece, the reader is referred to a recent compilation available in Bortolotti et al., (2013).

DISCUSSION AND CONCLUSIONS

Biosiliceous sedimentary rocks are widely known in the Dinarides, having originated in different geological settings throughout the Mesozoic (e.g., Vishnevskaya and Djerić, 2009; Vishnevskaya et al., 2009). They mostly appear as radiolarian cherts, both Triassic and Jurassic in age, found as blocks incorporated within the chaotic mélange sequences (e.g., Dimitrijević, 1997), and occasionally forming the interpillow matrix of the ophiolite extrusives (e.g., Halamić et al., 1999). The Jurassic radiolarian chert is also reported from non-ophiolite sequences of passive Adria margins where it stands as the ophiolitic mélange footwall (Vishnevskaya et al., 2009, Fig 2). Biostratigraphy makes an essential tool for revealing the age of sediments and,




Fig. 5 - Radiolarian age constraints in ophiolite complexes of the Central Dinaridic Ophiolite Belt. Data from the Mirdita Zone, Albania (Chiari et al., 2004), the western ophiolite belt in Greece (Chiari et al., 2003; Bortolotti et al., 2008; Nirta et al., 2010), Croatia (Halamčić et al. 1999), Serbia (Gawlick et al. 2009) and Bosnia (this study). Numerical ages are given according to Gradstein et al. (2012). The UAZ 95 column refers to Unitary Association Zones of Baumgartner et al. (1995b).

from this, deriving the age on thrusting events that have led to the ophiolite mélangé formation in the footwall of obducted ophiolites. Additionally, a variety of tectono-stratigraphic units is found incorporated in the mélangé, originated in different geological periods and settings, thus making mélangé a valuable source of information for the ocean evolution reconstruction.

Our data on radiolarians from the CDOB ophiolitic mélangé in Bosnia and Herzegovina point to their latest Bajocian to early Bathonian age. Assuming the formation of mélangé to be primarily a tectonic event, the reported time span thus documents the likely minimum ages of sedimentation of a siliceous sequence that made up the mélangé. In other words, the recovered ages do not necessarily coincide with the exact timing of the mélangé formation. A somewhat different line of reasoning is demonstrated by Gawlick et al. (2008), wherein the cherty sediments from an analogue ophiolite mélangé in northern Albania are considered to represent the matrix of an already disturbed oceanic sequence, thus conveniently marking the age of mélangé formation. As of yet, in the Dinarides, geographically the closest dated MOR-affiliated rocks from the ophiolite upper plate that tectonically overlies the analysed mélangé are olivine-bearing cumulate gabbros reported from the southern slopes of the Kozara Mountain, located about 50 km westwards from the sampling locality of Papratnica (Fig. 1; Ustaszewski et al., 2009). With the Sm-Nd age of approx. 158 ± 7.6 Ma, these gabbros stand for the youngest possible age of sea-floor spreading in this segment of the Neotethys. The latter thus effectively precludes the possibility that the mélangé formation was completed by mid-Jurassic times, but suggests it must have continued until after the MOR-type crust has ceased to generate which, given the possible analytical errors, might have lasted even until the Oxfordian-Kimmeridgian boundary. This stage was eventually completed with the main branch of the Dinaridic Neotethyan closure by the earliest Cretaceous at latest (Pamić et al., 2002), where only a few marginal back-arc basins remained opened thereafter (e.g., the Kozara

basin, Ustaszewski et al., 2009; Repno Oceanic Domain, Šegvić et al., in review). The age of MOR-type gabbros further corresponds to the K-Ar metamorphic ages of sub-ophiolitic soles recovered from the Krivaja-Konjuh ophiolite complex (161 ± 4 Ma; Lanphere et al., 1975) and the nearby Borja ophiolite complex (Fig. 1, 171.4 ± 3.7 Ma; Lugović et al., 2006), suggesting that the late ridge magmatism in the Dinaridic Neotethys was complemented by the near-ridge sole metamorphism, giving birth to the initial intraoceanic thrusting (e.g., Operta et al., 2003; Šegvić, 2010). Moreover, the established sole age denotes the earliest possible inception of the mélangé formation through the infill of the newly formed subduction trenches by gravitation that derived hangingwall material. In the Papratnica mélangé, fragments of metamorphic rocks are associated with mafic and ultramafic slide blocks that occur higher in the succession than the cherty sequence sampled for radiolarians (Fig. 2, Pamić, 1968). Another piece of evidence sheds light on the rapid evolution of the marginal Dinaridic Neotethys and the coeval mélangé formation. The pillow basalts containing radiolarian chert cropping out near the Ograjina River, situated a few kilometres away from the studied section, are proven to be of island-arc tholeiite (IAT) affinity (Simat, 2010). Analogue geochemistry is reported from the rest of the pillow basalts cropping out in the KKOC, and their origin is linked to advanced subduction processes and mantle wedge metasomatism, giving rise to IAT magmatism at ophiolite upper plate (Babajić, 2009). Given a subduction angle of 30° and an average subduction velocity of 5 cm *per annum* (for comparison the subduction of large Pacific plate is the fastest recent analogue, being subducted at the rate of 10 cm *per annum*, Dewey, 1981) will require approx. 3-4 Ma for the subducted material to attain a 110 km depth, given that most of the arc volcanism generates at this distance from the Benioff Zone (Murphy, 2006). In addition, the contribution from the subducted slab to island-arc lava sources appear to occur approximately 5 Ma prior to eruption (Turner, 2002), therefore making it no less than 10 Ma from the onset of intraoceanic subduction

(i.e. subophiolitic sole formation) to the first appearance of IAT lavas. The latter would also mean that the formation of ophiolitic mélangé in the Dinarides lasted at a minimum 10 Ma, encompassing both the primary sedimentation and the tectonic incorporation of the broad-provenanced olistoliths blocks.

A large grey carbonate olistolith, recognised in the Papratnica mélangé sequence, is by the analogy with the equivalent mapped carbonate blocks defined as Triassic in age (Fig. 2; Pamić, 1968). These limestones are apparently fully recrystallized, however, still featured by the preserved fossil remains that include the brachiopod macrofauna (*Rhynchonella vivida*, *Rhynchonella cf. mentzelli*, *Spirigera (Stolzenburgiella) bukowskii*, and *Spirigera (Pexidella) munster*), which along with the foraminifers (*Meandrospira dinarica*, *Frondicularia* sp., and *Pilamina densa*) indicate the Anisian pelagic to hemipelagic sedimentation (Pamić, 1968). Such peculiar Triassic components might be comparable to the so-called Hallstatt facies, usually comprising grey cherty limestones that can vary into red-bedded or light-coloured massive limestones known to have been deposited on the distal deeper shelf of the continental margin, being reworked and found in the Jurassic mélangé of the Northern Calcareous Alps in Austria (e.g., Mandl, 2000), but also in the contemporary mélangé of the Mirdita Zone in northern Albania (Gawlick et al., 2008), and elsewhere along the Middle Jurassic suture zone (Karamata et al., 1980; Robertson et al., 2013). This allows us to suggest that similar depositional conditions might have prevailed along the north-eastern portions of, at the time, the still integral Adriatic microcontinent (platform fragment of Gondwana), enabling the deposition of peculiar greyish Anisian limestones. Frontal to CDOB mélangé, a discontinuous radiolarite zone up to 800 m thick is situated, with a radiolarite sequence that covers a large stratigraphic interval from Middle Jurassic to Early Cretaceous (Pamić, 1982). The most complete section, Jezeračka Rijeka (locality no. 2 in Fig. 3), is composed of three lithostratigraphic units - varicoloured radiolarian chert (Bajocian to Oxfordian), red clayey chert and subordinately cherty limestone (Kimmeridgian to early Tithonian), and clayey cherty limestone with calpionellids (Tithonian - Berriasian) (Vishnevskaya and Djeric, 2009; Vishnevskaya et al., 2009). This succession of facies along with local enrichment in sponge spicules is indicative of continental slopes of the passive margin setting (Vishnevskaya et al., 2009). The fact that this sequence does not make part of the neighbouring mélangé is in favour of the hypothesis that, most likely, mainly the distal portions of the passive continental Triassic margin of the Adriatic microplate were dismembered and thereupon obliterated during the mélangé formation, preserved in mélangé in form of exotic limestone blocks of Triassic up to Tithonian age (Pamić, 1982).

Based on the above discussion, the following conclusions can be put forth:

1. The composite and chaotic nature of the material that numbers the fragments of the oceanic lithosphere and the associated sediments recovered from ophiolite mélangé of the CDOB (the Papratnica section in KKOC, NE Bosnia and Herzegovina) corroborates recent approaches advocating the mélangé formation in the ophiolite foot-wall wherein the radiolarite-ophiolite sequence originated from the obducted plate, whilst exotic limestone blocks must have provenanced from the Triassic continental margins.

2. he analysed radiolarian assemblage yielded the biostratigraphic age corresponding to the latest Bajocian to early Bathonian, being as such the first record on the age of radiolarians from an ophiolitic mélangé sequence from the Bosnian (central) portion of the CDOB.
3. Given the age of MOR-provenanced cumulate gabbros (158 ± 7.6 Ma) from the nearby upper-plate ophiolites of Mt. Kozara, known to be the youngest ocean ridge rocks of this segment of the Neotethys, we infer that the mélangé formation process could have lasted even until the Oxfordian-Kimmeridgian boundary (i.e. cessation of ocean spreading). Hence, the reported radiolarian age designates the time of sedimentation that might have preceded or, eventually, it coincided with the mélangé formation in this segment of the Dinarides.
4. The age of the metamorphic sole reported from the CDOB mélangé blocks (161 ± 4 Ma) overlaps with the crystallisation ages of MOR-affiliated gabbros. This has a two-fold signification: firstly, it denotes a rapid ocean setting transition, from an active ridge spreading to incipient intraoceanic subduction and, secondly, it stands for the earliest possible onset of the mélangé formation in the Dinarides.
5. IAT pillow basalts are also known from the mélangé and, referring to the beginning of intraoceanic subduction (i.e. sole metamorphism), it must have taken at least 10 Ma for arc-lavas to form. This further defines a minimum duration of thrusting tectonics that led to the genesis of the CDOB mélangé.
6. Anisian(?) limestones reported in the mélangé of the KKOC originally made part of the continental shelf of the Middle Triassic Adriatic microcontinent (to be disintegrated by Late Triassic into three separate entities, one being the Adria plate (e.g., Vlahović et al., 2005), suggesting that principally the distal shelf portions of the passive continental margin had been incorporated into the obducted mélangé, thus effectively constraining the magnitude of westward thrusting.
7. Final remarks on ophiolite thrusting in the Dinarides are in line with the fact that the ages of the upper-plate ophiolite suite - MOR gabbros (as young as 158 ± 7.6 Ma) and metamorphic sole (as young as 160 ± 4 Ma) - apparently mark younger ages (up to the Oxfordian) than the latest Bajocian to early Bathonian sedimentation ages of the ocean-floor pelagic cover, which made up the mélangé sampled for this study. The obduction tectonics (i.e. mélangé formation) was, therefore, not completed by the time indicated by the upper limit of radiolarian age (early Bathonian), but instead, it must have lasted at least until about the Oxfordian time. This corresponds well to the biostratigraphic data from the Serbian segment of the CDOB (Gawlick et al., 2009) and northern Albania (Chiari et al., 2004), where sediments forming the mélangé are as young as Callovian and early Oxfordian. This work also shows that biostratigraphy is an important tool for dating sediments and, from this, it serves as a venue to put constraints on thrusting events that caused the ophiolite mélangé formation. However, it does not by itself provide an age of mélangé, which is basically a tectonic event. The biostratigraphic dating therefore needs to be combined with a vigorous mélangé compositional analysis, principally of its crystalline components, in order to infer more on the time and palaeogeography of its genesis.

ACKNOWLEDGMENTS

Financial support for the field works in Bosnia and Herzegovina provided by the Faculty of Mining, Geology and Petroleum Engineering (University of Zagreb/Croatia) is gratefully acknowledged. The support from the research grant given by the Federal Ministry of Science and Culture of Bosnia and Herzegovina (03-39-7194-27/07) is appreciated. U. Barudžija (University of Zagreb/Croatia) is thanked for his assistance with petrographical investigations. We further extend our acknowledgment to Z. Jukić-Dupli (Žepče/Bosnia and Herzegovina) for his kind hospitality during the fieldwork. E. Gierlowski-Kordesch (Ohio University/USA) and A. Ristić are thanked for their help with English. Critical comments by K. Ustaszewski (Universität Jena/Germany) and Da. Slovenec (Croatian Geological Survey/Zagreb) greatly helped to improve an early version of the manuscript. Finally, constructive reviews by M. Chiari and one anonymous reviewer, as well as the editorial handling and suggestions by L. Pandolfi contributed significantly to the manuscript quality.

REFERENCES

- Aiello I.W., Hagstrum J.T. and Principi G., 2008. Peri-equatorial paleolatitudes for Jurassic radiolarian cherts of Greece. *Tectonophysics*, 448: 33-48.
- Aita Y., 1987. Middle Jurassic to Lower Cretaceous radiolarian biostratigraphy of Shikoku with reference to selected sections in Lombardy Basin and Sicily. *Sci. Reps. Tohoku Univ., Series 2 (Geology)*, 58: 1-91.
- Babajić E., 2009. Petrološko-geochemijska obilježja mafitnih stijena Krivajsko-konjuškog ofiolitnog kompleksa. Ph.D. thesis, Univ. Tuzla, Tuzla, Bosnia and Herzegovina, 158 pp.
- Baumgartner P.O., Bartolini A., Carter E.S., Conti M., Cortese G., Danelian T., De Wever P., Dumitrica P., Dumitrica-Jud R., Goričan Š., Guex J., Hull D.M., Kito N., Marcucci M., Matsuoka A., Murchev B., O'Dogherty L., Savary J., Vishnevskaya V., Widz D. and Yao A., 1995b. Middle Jurassic to Early Cretaceous radiolarian biochronology of Tethys based on Unitary Associations. In: P.O. Baumgartner, L. O'Dogherty, Š. Goričan, E. Urquhart, A. Pillevuit and P. De Wever (Eds.), *Middle Jurassic to Lower Cretaceous Radiolaria of Tethys: Occurrences, systematics, biochronology*. *Mém. Géol., Lausanne*, 23: 1013-1048.
- Baumgartner P.O., O'Dogherty L., Goričan Š., Dumitrica-Jud R., Dumitrica P., Pillevuit A., Urquhart E., Matsuoka A., Danelian T., Bartolini A., Carter E.S., De Wever P., Kito N., Marcucci M. and Steiger T., 1995a. Radiolarian catalogue and systematics of Middle Jurassic to Early Cretaceous Tethyan genera and species. In: P.O. Baumgartner, L. O'Dogherty, Š. Goričan, E. Urquhart, A. Pillevuit and P. De Wever (Eds.), *Middle Jurassic to Lower Cretaceous Radiolaria of Tethys: Occurrences, systematics, biochronology*. *Mém. Géol., Lausanne*, 23: 37-685.
- Bortolotti V., Chiari M., Marcucci M., Marroni M., Pandolfi L., Principi G. and Saccani E., 2004. Comparison among the Albanian and Greek ophiolites: in search of constraints for the evolution of the Mesozoic Tethys Ocean. *Ophioliti*, 29: 19-35.
- Bortolotti V., Chiari M., Marcucci M., Photiades A., Principi G. and Saccani E., 2008. New geochemical and age data on the ophiolites from the Othrys area (Greece): implication for the Triassic evolution of the Vardar Ocean. *Ophioliti*, 33: 135-151.
- Bortolotti V., Chiari M., Marroni M., Pandolfi L., Principi G. and Saccani E., 2013. Geodynamic evolution of ophiolites from Albania and Greece (Dinaric-Hellenic belt): one, two, or more oceanic basins? *Intern. J. Earth Sci.*, 102: 783-811.
- Bortolotti V., Kodra A., Marroni M., Mustafa F., Pandolfi L., Principi G. and Saccani E., 1996. Geology and petrology of ophiolitic sequences in the Mirdita region (Northern Albania). *Ophioliti*, 21: 3-20.
- Chiari M., Bortolotti V., Marcucci M., Photiades A. and Principi G., 2003. The Middle Jurassic siliceous sedimentary cover at the top of the Vourinos Ophiolite (Greece). *Ophioliti*, 28: 95-103.
- Chiari M., Bortolotti V., Marcucci M., Photiades A., Principi G. and Saccani E., 2012. Radiolarian biostratigraphy and geochemistry of the Koziakas massif ophiolites (Greece). *Bull. Soc. Géol. France*, 183: 287-306.
- Chiari M., Djerić N., Garfagnoli F., Hrvatović H., Krstić M., Levi N., Malasoma A., Marroni M., Menna F., Nirta G., Pandolfi L., Principi G., Saccani E., Stojadinović U. and Trivić B., 2011. The geology of the Zlatibor-Maljen area (western Serbia): a geotraverse across the ophiolites of the Dinaric-Hellenic collisional belt. *Ophioliti*, 36: 139-166.
- Chiari M., Marcucci M. and Prella M., 1994. Mirdita Ophiolites project: 2 Radiolarian assemblages in the cherts at Fusche Arrez and Shebaj (Mirdita Area, Albania). *Ophioliti*, 19: 313-318.
- Chiari M., Marcucci M. and Prella M., 2004. Radiolarian assemblages from the Jurassic cherts of Albania: new data. *Ophioliti*, 29: 95-105.
- Çollaku A, Cadet JP, Bonneau M and Jolivet L., 1992. L'édifice structural de l'Albanie septentrionale: des éléments de réponse sur les modalités de la mise en place des ophiolites. *Bull. Soc. Géol. France*, 163: 455-468.
- Danelian T. and Robertson A.H.F., 2001. Neotethyan evolution of eastern Greece (Pagondas Mélange, Evia island) inferred from radiolarian biostratigraphy and the geochemistry of associated extrusive rocks. *Geol. Mag.*, 138: 345-363.
- Dewey J.F., 1981. Episodicity, sequence and style at convergent plate boundaries. In: *The continental crust and its mineral deposits*. *Geol. Ass. Can., Spec. Paper*, 20: 553-572.
- Dimitrijević M.D., 1997. *Geology of Yugoslavia*. *Geol. Inst. GEMINI, Belgrade*, 187 pp.
- Dimitrijević M.N., Dimitrijević M.D., Karamata S., Sudar M., Gerzina N., Kovács S., Dosztály L., Gulácsi Z., Less Gy. and Pelikán P., 2003. Olistrotrome/mélanges - an overview of the problems and preliminary comparison of such formations in Yugoslavia and NE Hungary. *Slovak Geol. Mag.*, 9: 3-21.
- Dragičević I., 1987. Paleogeografska evolucija rubnog dijela mezozojskog karbonatnog šelfa Dinarida između Vrbasa i Bosne. Ph.D. thesis, Univ. Zagreb, Zagreb, Croatia, 90 pp.
- Gawlick H.J., Frisch W., Hoxha L., Dumitrica P., Krystyn L., Lein R., Missoni S. and Schlagintweit F., 2008. Mirdita Zone ophiolites and associated sediments in Albania reveal Neotethys Ocean origin. *Intern. J. Earth Sci.*, 97: 865-881.
- Gawlick H.J., Sudar M., Suzuki H., Đerić N., Missoni S., Lein R. and Jovanović D., 2009. Upper Triassic and Middle Jurassic radiolarians from the ophiolitic mélange of the Dinaridic Ophiolite Belt, SW Serbia. *N. Jahrb. Geol. Paläont. Abhandl.*, 253: 293-311.
- Gradstein F.M., Ogg J.G., Schmitz M.D. and Ogg G.M. 2012. *The Geological Time Scale 2012*. Elsevier, Amsterdam, 1176 pp.
- Halamić J., Goričan, Š., Slovenec, D. and Kolar-Jurkovšek, T., 1999. A Middle Jurassic radiolarite-clastic succession from the Medvednica Mt. (NW Croatia). *Geol. Croatica*, 52: 29-57.
- Hrvatović H., 2006. *Geološki vodič kroz Bosnu i Hercegovinu*, Geol. Survey of Federation of Bosnia and Herzegovina, Ilidza (Sarajevo), 203 pp.
- Jovanović R., Mojičević M., Tokić S. and Rokić Lj., 1978. Petrološki izvještaj za tumač lista Sarajevo, 1:100000 osnovne geološke karte SFRJ. *Geol. Surv. Sarajevo - Federal Geol. Surv., Belgrade*.
- Karamata S., 2006. The geodynamical framework of the Balkan Peninsula: its origin due to the approach, collision and compression of Gondwanian and Eurasian units. In: A.H.F. Robertson and D. Mountrakis (Eds.), *Tectonic development of the Eastern Mediterranean Region*. *Geol. Soc. London Spec. Publ.*, 260: 155-178.
- Karamata S., Majer V. and Pamić J., 1980. The ophiolites of Yugoslavia. In: G. Rocci (Ed). *Tethyan ophiolites*. *Ophioliti*, special issue, vol. 1, Florence, p. 170-189.
- Kellici I., de Wever P. and Kodra A., 1994. Radiolaires mésozoïques du massif ophiolitique de Mirdita, Albanie. *Paléontologie et stratigraphie*. *Rev. Micropal.*, 37: 209-222.

- Kolodny Y., Chaussidon M. and Katz A., 2005. Geochemistry of a chert breccia. *Geochem. Cosmochem. Acta*, 69: 427-439.
- Lanphere M., Coleman R.C., Karamata S. and Pamić J., 1975. Age of amphibolites associated with alpine peridotites in the Central Dinaride Ophiolite Belt, Yugoslavia. *Earth Planet. Sci. Lett.*, 26: 271-276.
- Lugović B., Altherr R., Raczek I., Hofmann A.W. and Majer V., 1991. Geochemistry of peridotites and mafic igneous rocks from the Central Dinaridic Belt, Yugoslavia. *Contrib. Mineral. Petrol.*, 106: 201-216.
- Lugović B., Šegvić B., Babajić E. and Trubelja F. 2006. Evidence of short-living intraoceanic subduction in the Central Dinarides, Konjuh ophiolite complex (Bosnia-Herzegovina). In: K. Resim-ić-Šarić (Ed.). *Mesozoic ophiolite belts of northern part of the Balkan Peninsula. Proceed. Univ. Belgrade, Faculty Mining and Geology, Belgrade*, p. 72-75.
- Majer V., Kreuzer J., Harre W., Seidel E., Altherr R and Okrusch M., 1979. Petrology and geochronology of metamorphic rocks from Banija, Yugoslavia. *Intern. Ophiolite. Symp. Abstr. (Nicosia)*: 46-47.
- Mandl G.W., 2000. The Alpine sector of the Tethyan shelf - Examples of Triassic to Jurassic sedimentation and deformation from the Northern Calcareous Alps. *Mitt. Österr. Geol. Gesell.*, 92: 61-77.
- Marcucci M., Kodra A., Pirdeni A. and Gjata T., 1994. Radiolarian assemblages in the Triassic and Jurassic cherts of Albania. *Ofioliti*, 19: 105-114.
- Murphy J.B., 2006. Igneous rock associations. 7. Arc magmatism, I: Relationship between subduction and magma genesis. *Geosci. Can.*, 33 (4): 145-167.
- Nirta G., Bortolotti V., Chiari M., Menna F., Saccani E., Principi G. and Vannucchi P., 2010. Ophiolites from the Grammos-Arrenes area, northern Greece: geological, paleontological and geochemical data. *Ofioliti*, 35: 103-115.
- O'Dogherty L., Bill M., Goričan Š., Dumitrica P. and Masson H. 2006. Bathonian radiolarians from an ophiolitic mélange of the Alpine Tethys (Gets Nappe, Swiss-French Alps). *Micropal.*, 51: 425-485.
- O'Dogherty L., Carter E.S., Dumitrica P., Goričan Š., De Wever P., Bandini A.N., Baumgartner P.O. and Matsuoka A., 2009. Catalogue of Mesozoic radiolarian genera. Part 2: Jurassic - Cretaceous. In: L. O'Dogherty, Š. Goričan and P. De Wever (Eds.), *Catalogue of Mesozoic radiolarian genera. Geodiversitas*, 31: 271-356.
- Operta M., Pamić J., Balen D. and Tropper P., 2003. Corundum-bearing amphibolites from the metamorphic basement of the Krivaja-Konjuh ultramafic massif (Central Dinaride Ophiolite Belt, Bosnia). *Mineral. Petrol.*, 77: 287-295.
- Pamić J., 1968. Petrološki izvještaj za tumač lista Zavidovići, 1:100000 osnovne geološke karte SFRJ. *Geol. Surv. Sarajevo - Federal Geol. Surv. Belgrade*.
- Pamić J., 1970. Petrološki izvještaj za tumač lista Vareš, 1:100000 osnovne geološke karte SFRJ. *Geol. Surv. Sarajevo - Federal Geol. Surv. Belgrade*.
- Pamić J., 1982. Some geological problems of the Dinaridic ophiolites and their associations. *Earth Evol. Sci.*, 2: 30-35.
- Pamić J. and Hrvatović H., 2000. Dinaride Ophiolite Zone (DOZ). Basic data on the geology and petrology of the Krivaja-Konjuh ophiolite complex. In: J. Pamić and B. Tomljenović (Eds.), *Pancardi 2000 fieldtrip guidebook. Vijesti Hrvatskoga geološkog društva*, 37 (2): 60-68.
- Pamić J., Gušić I. and Jelaska V., 1998. Geodynamic evolution of the central Dinarides. *Tectonophysics*, 297: 251-268.
- Pamić J., Sunarić O. and Olujčić J., 1979. Tumač osnovne geološke karte SFRJ, list Teslić. *Geol. Surv. Sarajevo - Federal Geol. Surv. Belgrade*.
- Pamić J., Sunarić-Pamić O., Olujčić J. and Antić R., 1977. Petrografija i petrologija krivajsko-konjuškog ofiolitskog kompleksa i njegove osnovne geološke karakteristike. *Acta Geol. (Zagreb)*, 9: 59-135.
- Pamić J., Tomljenović B. and Balen D., 2002. Geodynamic and petrogenetic evolution of Alpine ophiolites from the central and NW Dinarides: an overview. *Lithos*, 65: 113-142.
- Prela M., Chiari M. and Marucci M., 2000. Jurassic radiolarian biostratigraphy of the sedimentary cover of ophiolites in the Mirdita Area, Albania: new data. *Ofioliti*, 25: 55-62.
- Robertson A.H.R., 2002. Overview of the genesis and emplacement of Mesozoic ophiolites in the Eastern Mediterranean Tethyan region. *Lithos*, 65: 1-67.
- Robertson A.H.R., Trivić B., Đerić N and Bucur I., 2013. Tectonic development of the Vardar Ocean and its margins: Evidence from the Republic of Macedonia and Greek Macedonia. *Tectonophysics*, 595-596: 25-54.
- Schmid S.M., Bernoulli D., Fügenschuh B., Matenco L., Schefer S., Schuster R., Tischler M. and Ustaszewski K., 2008. The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units. *Swiss J. Geosci.*, 101:139-183.
- Šegvić B., 2010. Petrologic and geochemical characteristics of the Krivaja-Konjuh ophiolite complex (NE Bosnia) - petrogenesis and regional geodynamic implications. Ph.D. thesis, Univ. Heidelberg, Heidelberg, Germany, 301 pp.
- Šegvić B., Lugović B., Slovenec D., Meyer H-P. and Ignjatić S. in review. Lower Cretaceous metamorphic sole from the Kalnik Mt. (North Croatia): Evidence of initial convergence in a back-arc marginal basin of Meliata-Maliac - Dinaric-Vardar oceanic systems. *Austrian J. Earth Sci.*, in review.
- Simat S., 2010. Geokemijske i tektonomagmatske karakteristike jastučastih bazalta iz ofiolitnih melanza Dinarida. Master thesis, Univ. Zagreb, Zagreb, Croatia, 66 pp.
- Smith A.G. and Spray J.G., 1984. A half-ridge transform model for the Hellenic-Dinaric ophiolites. In: J.E. Dixon and A.H.F. Robertson (Eds.), *The geological evolution of the Eastern Mediterranean. Geol. Soc. London Spec. Publ.*, 17: 629-644.
- Spray J.G., Bébien J., Rex D.C. and Roddick J.C., 1984. Age constraints on the igneous and metamorphic evolution of the Hellenic-Dinarides ophiolites. In: J.E. Dixon and A.H.F. Robertson (Eds.), *The geological evolution of the Eastern Mediterranean. Geol. Soc. London Spec. Publ.*, 17: 619-627.
- Tari V., 2002. Evolution of the northern and western Dinarides: a tectonostratigraphic approach. In: G. Bertotti, K. Schulmann and S. Cloetingh (Eds.), *Continental collision and the tectono-sedimentary evolution of forelands. Copernicus, Katlenburg-Lindau*, p. 223-236.
- Turner S.P., 2002. On the time-scales of magmatism at island-arc volcanoes. *Philos Trans A Math Phys Eng Sci.*, 360: 2853-2871.
- Ustaszewski K., Schmid S., Lugović B., Schuster R., Schaltegger U., Bernoulli D., Hottinger L., Kounov A., Fügenschuh B. and Schefer S., 2009. Late Cretaceous intra-oceanic magmatism in the internal Dinarides (northern Bosnia and Herzegovina): Implications for the collision of the Adriatic and European plates. *Lithos*, 108: 106-125.
- Ustaszewski K., Kounov A., Schmid S., Schaltegger U., Krenn E., Frank W. and Fügenschuh B., 2010. Evolution of the Adria-Europe plate boundary in the northern Dinarides: From continent-continent collision to back-arc extension. *Tectonics*, 29: 1-34.
- Vishnevskaya V.S. and Djerić N. 2009. Mesozoic radiolaria of Bosnia and Serbia: New data. *Paleont. J.*, 43: 1513-1568.
- Vishnevskaya V.S., Djerić N. and Zakariadze G.S., 2009. New data on Mesozoic radiolaria of Serbia and Bosnia, and implications for the age and evolution of oceanic volcanic rocks in the Central and Northern Balkans. *Lithos*, 108: 72-105.
- Vlahović I., Tišljarić J., Velić I. and Matičec D., 2005. Evolution of the Adriatic carbonate platform: Paleogeography, main events and depositional dynamics. *Palaeo. Palaeo. Palaeo.*, 220: 333-360.
- Yao A., 1979. Radiolarian fauna from the Mino Belt in the northern part of the Inuyama Area, Central Japan, Part II: Nassellaria, 1. *J. Geosci., Osaka City Univ.*, 22: 21-72.




Plate 1 - Middle Jurassic radiolarians from the analysed sample. For each specimen, the scanning electron microscopograph number is indicated. The rock sample, residue and illustrated specimens are stored at the Ivan Rakovec Institute of Paleontology, ZRC SAZU, Ljubljana, Slovenia. 1) *Archaeodictyomitra rigida* Pessagno, 100233. This specimen has less costae and larger pores than the type material but it fits well in the variability of *Archaeodictyomitra rigida* as considered by O'Dogherty et al. (2006); 2, 3) *Archaeodictyomitra suzukii* Aita, 2- 100114, 3- 100124; 4) *Archaeodictyomitra* sp., 100230; 5, 6) *Semihsum amabile* (Aita), 5- 100132, 6- 100115; 7) *Dictyomitrella* (?) *kamoensis* Mizutani and Kido, 100150; 8, 9) *Canoptum krahsteinense* (Suzuki and Gawlick), 8- 100226, 9- 100227; 10) *Eoxitus baloghi* Kozur, 100101; 11) *Transhsuum maxwelli* (Pessagno) gr., 100134; 12) *Transhsuum brevicostatum* (Ožvoldova) gr., 100133; 13) *Stichomitra* (?) *takanoensis* Aita gr., 100228; 14) *Stichocapsa* (?) *magnipora* Chiari, Marcucci and Prela, 100104; 15) *Eucyrtidiellum semifactum* Nagai and Mizutani, 100105. This specimen has longer plicae than the holotype but it conforms well with other specimens included in the variability of *Eucyrtidiellum semifactum* (see Baumgartner et al., 1995a, pl. 3016, fig. 3(H) and Figs. 1, 2, respectively); 16) *Eucyrtidiellum unumaense dentatum* Baumgartner, 100149; 17) *Quarticella* sp., 100126; 18) *Triactoma* (?) sp., 100235; 19) *Unuma laticostatus* (Aita), 100116; 20) *Yaocapsa* sp. A, 100236. The species is close to *Yaocapsa* sp. aff. *Y. mastoidea* (Yao) (it was *Stichocapsa* sp. E in Baumgartner et al., 1995a, p. 524, pl. 4042, figs. 1, 2) but differs from the latter by having faint ridges that connect the adjacent pores; 21) *Praewilliriedellum* sp. aff. *P. convexum* (Yao), 100124. The species differs from the type material (in Yao, 1979) by having a higher and wider abdomen; 22) *Praewilliriedellum* sp. aff. *P. himedaruma* (Aita), 100237. The species differs from the type material (in Aita, 1987) by having abdomen narrower than the last segment; 23) *Protunuma lanosus* Ožvoldova, 100117; 24) *Paronaella broennimanni* Pessagno, 100229; 25a-b) *Unuma* sp., 25a- 100122, 25b- antapical view, 100121; 26a-b) *Unuma gordus* Hull, 26a- 100231, 26b- antapical view, 100232; 27a-b) *Striatojaponocapsa synconexa* O'Dogherty, Goričan and Dumitrica, 27a- 100154, 27b- antapical view, 100155; 28a-b) *Hemicryptocapsa marcucciae* (Cortese), 28a- 100138, 28b- antapical view, 100137; 29a-b) *Hemicryptocapsa yaoi* (Kozur), 29a- 100110, 29b- antapical view, 100111; 30a-b) *Helvetocapsa matsukoi* (Sashida), 30a- 100239, 30b- antapical view, 100238.

