

# Annals of Clinical Biochemistry: An international journal of biochemistry and laboratory medicine

<http://acb.sagepub.com/>

---

**Oxidative status and the severity of clinical symptoms in patients with post-traumatic stress disorder**  
Leda Borovac Stefanovic, Dubravka Kalinic, Ninoslav Mimica, Blanka Beer Ljubic, Jasna Aladrovic, Marina Mandelsamen  
Perica, Maja Curic, Petra Folnegovic Grosic and Ivancica Delas  
*Ann Clin Biochem* published online 4 April 2014  
DOI: 10.1177/0004563214528882

The online version of this article can be found at:  
<http://acb.sagepub.com/content/early/2014/03/31/0004563214528882>

---

Published by:



<http://www.sagepublications.com>

On behalf of:

Association for Clinical Biochemistry and Laboratory Medicine



The Association for  
**Clinical Biochemistry &  
Laboratory Medicine**

Netherlands Society for Clinical Chemistry and Laboratory  
Japan Society of Clinical Chemistry

Additional services and information for *Annals of Clinical Biochemistry: An international journal of biochemistry and laboratory medicine*  
can be found at:

**Email Alerts:** <http://acb.sagepub.com/cgi/alerts>

**Subscriptions:** <http://acb.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

>> [OnlineFirst Version of Record](#) - Apr 4, 2014

[What is This?](#)



Annals of Clinical Biochemistry  
0(0) 1–10  
© The Author(s) 2014  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/0004563214528882  
acb.sagepub.com



## Oxidative status and the severity of clinical symptoms in patients with post-traumatic stress disorder

Leda Borovac Štefanović<sup>1</sup>, Dubravka Kalinić<sup>1,2</sup>, Ninoslav Mimica<sup>1,2</sup>, Blanka Beer Ljubić<sup>3</sup>, Jasna Aladrović<sup>3</sup>, Marina Mandelsamen Perica<sup>1</sup>, Maja Ćurić<sup>1</sup>, Petra Folnegović Grošić<sup>4</sup> and Ivančica Delaš<sup>1</sup>

### Abstract

**Background:** The aim of this study was to measure the parameters of oxidative stress in the blood of patients with post-traumatic stress disorder.

**Methods:** The study included 80 male war veterans who participated actively in the Homeland war in Croatia. Volunteers were divided into two groups: 50 veterans diagnosed with post-traumatic stress disorder and 30 without diagnosis. The self-assessment Hospital Anxiety and Depression Scale and the Beck Depression Inventory were used to detect the severity of depression and anxiety in the post-traumatic stress disorder patients. Catalytic concentrations of superoxide dismutase and glutathione peroxidase in erythrocytes and the concentration of malondialdehyde in serum were measured spectrophotometrically.

**Results:** Although the catalytic concentrations of erythrocyte superoxide dismutase and erythrocyte glutathione peroxidase were within the reference range for both groups, the values obtained for the post-traumatic stress disorder group were significantly lower ( $P < 0.001$ ). For serum malondialdehyde concentrations, no statistically significant differences between the groups were found.

**Conclusions:** Lower catalytic concentrations of erythrocyte superoxide dismutase and erythrocyte glutathione peroxidase in patients with post-traumatic stress disorder may indicate a weaker response to oxidative stress due to impaired enzyme activity and/or decreased synthesis. Conversely, no significant changes in serum malondialdehyde concentrations suggest a compensated balance and adaptive response to (oxidative) stress.

### Keywords

Enzymes, nutrition, enzymatic methods

Accepted: 28th February 2014

### Introduction

Free radicals, atoms, molecules or ions with an unpaired number of electrons are physiological products of metabolism which directly participate in the protection against infection, or act as coordinators of inflammatory response.<sup>1</sup> In a healthy organism, the production and various interactions of free radicals are tightly controlled by enzymatic defence mechanisms

<sup>1</sup>School of Medicine, University of Zagreb, Zagreb, Croatia

<sup>2</sup>University Psychiatric Hospital Vrapče, Zagreb, Croatia

<sup>3</sup>Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

<sup>4</sup>Department of Psychiatry, University Hospital Centre Zagreb, Zagreb, Croatia

#### Corresponding author:

Ivančica Delaš, University of Zagreb, School of Medicine, Department of Chemistry and Biochemistry, Šalata 3, HR-10 000 Zagreb, Croatia.  
Email: ivancica.delas@mef.hr

such as the superoxide dismutase (SOD), glutathione peroxidase (GPx) and catalase (CAT), or via the action of non-enzymatic antioxidants such as ascorbic acid, Vitamin E and glutathione.<sup>2</sup> However, excessive production of free radicals and/or impaired protection mechanisms lead to oxidative stress and result in radical-induced damage of lipids, proteins and DNA.<sup>3</sup> Tissue damage caused by oxidative stress includes a diversity of biological phenomena such as aging, carcinogenesis, atherosclerosis and neurodegeneration.<sup>4</sup> The severity of oxidative stress can be measured via the concentration of reactive oxygen/nitrogen species (ROS/RNS) or the concentration of antioxidants. Furthermore, biomolecules derived through the action of free radicals can be used as markers of oxidative stress: damaged DNA/RNA (i.e. 8-hydroxy(deoxy)guanosine), the oxidation/nitration of proteins (i.e. protein carbonyl content) and the products of lipid peroxidation (i.e. 4-hydroxy-2-nonenal, HNE; malondialdehyde, MDA).

Lipid peroxidation, defined as the oxidative deterioration of lipids containing any number of carbon-carbon double bonds, is considered to participate significantly in the creation of oxidative stress, since polyunsaturated fatty acids (PUFAs) frequently serve as a target for free radicals resulting in their multiplication through a cascade of reactions.<sup>1</sup> The primary products of lipid peroxidation, which include conjugated dienes and hydroperoxides, are relatively unstable and decompose into secondary products such as alcohols, aldehydes and ketones. Although numerous state-of-the-art methods have been developed, the concentration of overall lipid peroxidation is still most frequently assessed as the concentration of serum malondialdehyde (sMDA) which is determined by a reaction with thiobarbituric acid (TBARS Assay).

Lipids comprise about 60% of the brain's dry weight with a high ratio of PUFAs, and the brain utilizes more than 20% of total O<sub>2</sub> consumed by the body, which is why the brain is highly susceptible to lipid peroxidation that might result in serious damages and loss of function.<sup>5-7</sup> Oxidative stress may alter neurotransmission patterns, neuronal function and brain activity in general, and can also disrupt the membrane integrity.<sup>8,9</sup> Both the importance of fatty acid composition in brain cell membranes and the involvement of PUFAs in the development of various psychiatric disorders have been intensively studied.<sup>10-12</sup> A significant number of these results point to a relationship between the lack of essential PUFAs, particularly eicosapentaenoic (EPA, C20:5n-3) and docosahexaenoic (DHA, C22:6n-3), and a variety of psychiatric symptoms and disorders such as stress, anxiety, cognitive impairment, mood disorders and schizophrenia. In our previous study,<sup>13</sup> a significant negative correlation between

plasma EPA concentration and the severity of clinical symptoms in patients with post-traumatic stress disorder (PTSD) was found.

According to recently released DSM V criteria,<sup>14</sup> PTSD is classified as a trauma- and stressor-related disorder caused by a traumatic experience such as exposure to actual or threatened death, serious injury or sexual violation. It is characterized by four major symptom clusters: re-experiencing, arousal, avoidance and persistent negative alterations in cognitions and mood. The involvement of PTSD results in impairments in the individual's social interactions, work capability or everyday functioning, although it is unclear why certain individuals develop PTSD as a reaction to traumatic incidents while others do not.

Since oxidative stress is a general phenomenon and the markers of oxidative damage in the central nervous system correlate well with those in peripheral materials, the majority of analyses were performed on easily accessible materials such as blood, saliva and cerebrospinal fluid.<sup>5</sup> A substantial number of authors found increased amounts of markers of oxidative stress in plasma/serum, and/or deviations in the activity of defence enzymes in patients suffering from psychiatric disorders.<sup>3,7,15-19</sup> A recent review by Hovatta *et al.*<sup>20</sup> analysed available data on the relationship between oxidative stress and anxiety, depression and comorbid disorders, and the possible mechanisms of impaired defence against oxidative stress. Stress-induced alterations in the concentration of neurotransmitters such as excitatory amino acids, serotonin and gamma-amino-butyric acid can modulate the immune response and *vice versa*, and cytokines, mediators produced by immune cells, maintain homeostasis by communicating with other biological systems such as the hypothalamic-pituitary-adrenal (HPA) axis.<sup>21,22</sup>

Recently, Kuloglu *et al.*<sup>23,24</sup> recognized a link between oxidative stress and some anxiety disorders (obsessive-compulsive and panic disorder), demonstrating that oxidative metabolism can influence anxiety regulation. Bouayed *et al.*<sup>25</sup> in their review summarized data which demonstrate a link between oxidative stress and anxiety, but an unambiguous cause-effect relationship has yet to be determined. Some of the reviewed studies suggest oxidative stress causes anxiety-related behaviours but none clarify the underlying, specific mechanisms in detail.

The risk factors for developing PTSD appear to be related to a combination of genetic predisposition and environmental factors.<sup>26,27</sup> It was first recognized among Vietnam War veterans but was also subsequently diagnosed in people expressing similar symptoms in response to various traumas, whether in war or civil life.<sup>28</sup> During the 1990s, the Homeland War struck Croatia. This unfortunate experience resulted in a vast

number of individuals suffering from PTSD. Since PTSD has some clinical symptoms similar to those found in depression, bipolar disorder, schizophrenia and other psychiatric disorders, the present study investigates whether similar changes in oxidative/antioxidative balance could be confirmed. Certain lifestyle and nutritional habits, which are possible cofactors in the processes of oxidative stress, were also observed in the study participants.

## Subjects and methods

### Subjects

The study included 80 male Croatian Homeland War veterans. Fifty veterans were outpatients from University Psychiatric Hospital Vrapče who were diagnosed with PTSD according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV),<sup>29</sup> and the International Statistical Classification of Diseases and Related Health Problems (ICD-10)<sup>30</sup> diagnostic criteria. The control group included 30 participants who had also served in the Homeland War, but did not develop PTSD. The diagnosis evaluation was performed by experienced psychiatrists in order to confirm the PTSD diagnoses according to the DSM-IV and ICD-10. Although an updated edition of the manual (DSM-V)<sup>14</sup> was released in the meantime, where PTSD is detached from anxiety disorders and included in a new section of trauma- and stressor-related disorders, all our patients met the criteria for PTSD diagnosis according to both DSM IV and V. The self-assessment Hospital Anxiety and Depression Scale (HADS)<sup>31</sup> and the Beck Depression Inventory (BDI-II)<sup>32</sup> were used to detect the severity of depression and anxiety in the PTSD patients. The 21 BDI-II items, as well as 14 items from the HADS, are each scored on a four-point scale from 0 = 'no symptoms' to 3 = 'extreme symptoms'. The sum of the first seven items from the HADS is the severity index for anxiety, and the sum of the last seven is the index for depression. The severity of PTSD was evaluated by a clinician-rated instrument, i.e. the Clinical Global Impression – Severity scale (CGI-S).<sup>33</sup> The severity of the patient's PTSD is assessed by CGI-S in a seven point range: from 1 ('reference, not at all ill') to 7 ('extremely ill'). All the participants were free of all medication for at least two weeks prior to entering the study, with the exception of veterans with PTSD who were treated with an antidepressant (sertraline, 50 mg/day) for at least two months. Ten of the PTSD patients were also on therapy with anxiolytics (4 – diazepam; 3 – alprazolam; 3 – lorazepam). Exclusion criteria were: alcohol and substance abuse or dependence, food supplementation (i.e. vitamins E and/or C), presence of a severe neurologic

disorder, presence of an infectious disease or presence of a severe physical disorder. The study protocol was approved by the Ethics Committees of the University Psychiatric Hospital Vrapče and the School of Medicine University of Zagreb, Croatia. Prior to entering the study all participants signed an informed consent, and afterwards completed a general health questionnaire.<sup>34</sup>

### Blood sample collection

Vein blood samples were collected after overnight fasting into 5 mL vacutainer tubes (Becton Dickinson, Franklin Lakes, NJ, USA) containing ethylenediaminetetraacetic acid. The samples were then centrifuged at 3000 rpm for 10 min at 15°C to separate the plasma from blood cells. After washing procedures, erythrocytes isolated from blood samples were haemolysed by the freezing and thawing sequence.

### Measurement of antioxidant enzymes activity

Haemolysates of erythrocytes were used to measure catalytic concentrations of erythrocyte glutathione peroxidase (eGPx) (EC 1.11.1.9) and erythrocyte superoxide dismutase (eSOD) (EC 1.15.1.1), as well as haemoglobin (Hb) concentration. The eGPx activity in erythrocyte haemolysates was measured using a commercial RANSEL kit and eSOD activity was measured with RANSOD reagent (both from Randox Laboratories Ltd., Dublin, Ireland) on SABA 18 auto-analyzer (AMS, Rome, Italy). The eGPx activity, expressed in nanomoles of nicotinamide adenine dinucleotide phosphate oxidized per minute, was measured by a coupled modification of the assay procedure of Paglia and Valentine.<sup>35</sup> The assay for eSOD activity was based on the method described by Woolliams *et al.*,<sup>36</sup> using xanthine and xanthine oxidase to produce superoxide radicals which then react with nitroblue tetrazolium to form a red formazan dye. The inhibitory effects of eSOD on the initial rate of this reaction were measured, and one unit of SOD activity is that which causes a 50% inhibition of the rate of reduction of the substrate under the conditions of the assay. In haemolysates, a mass concentration of haemoglobin was also measured (Herbos Dijagnostika, Sisak, Croatia) and eGPx and eSOD catalytic concentrations were expressed in terms of U/g Hb.

### Determining sMDA concentrations

Serum MDA concentrations were measured using the modified TBARS method described by Anguloa *et al.*<sup>37</sup> and Botsoglou *et al.*<sup>38</sup> The results were calculated and

expressed as concentration of MDA ( $\mu\text{mol/L}$ ) using the extinction coefficient of  $156,000 \text{ Lmol}^{-1} \text{ cm}^{-1}$ .<sup>39</sup>

### Laboratory methods

Serum triacylglycerols (TG), total cholesterol (T-CH) and high-density lipoprotein cholesterol (HDL-CH) were measured by enzymatic colorimetric methods (Roche Diagnostics Corporation, Indianapolis, IN, US). Low-density lipoprotein cholesterol (LDL-CH) was calculated as described by Friedewald *et al.*<sup>40</sup>

### Statistical analysis

The obtained data were evaluated with STATISTICA 10 software (StatSoft, Inc., Tulsa, Oklahoma, USA). The data distribution was tested with Kolmogorov–Smirnov and Shapiro–Wilk’s W test. Descriptive statistics were performed, and the results are presented as mean  $\pm$  standard deviation (mean  $\pm$  SD). The results were processed by Student’s *t*-test, Mann–Whitney and Chi-square test, and the correlation of study parameters was tested by Spearman rank order correlations. The observed differences were considered statistically significant at the level of  $P < 0.05$ .

## Results

### Demographic data

A total of 80 male volunteers who participated actively in the Croatian Homeland War in the 1990s were enrolled in this study. The PTSD group included 50 outpatients with a mean age of 44.6 years (range 34–58 years). The sociodemographic and biochemical characteristics of the PTSD patients and the control group are summarized in Table 1. The participants from the PTSD group and control group did not differ significantly regarding the duration of exposure to combat trauma, nor in the time elapsed after the war trauma terminated. The degree of education was significantly higher within the control group (66.67% with higher education) while participants from the PTSD group had mainly a secondary school degree (81.63%). At the time of their participation in the study, most of the PTSD veterans were either retired or unemployed.

### Lifestyle and nutritional habits

Data on nutritional habits included information about the body mass index (BMI) as well as the consumption of fish and various oils (Table 1). The mean BMI value was similar for both groups. Regarding the oil type, sunflower oil was preferred in both groups with significantly higher consumption frequencies recorded in

patients with PTSD ( $P = 0.0203$ ). However, the control group consumed significantly more olive oil compared to the PTSD group ( $P = 0.0025$ ). For fish consumption, no significant differences were found between the groups. Analysis of laboratory data revealed no significant differences in lipid parameters between the two groups.

### Severity of depression and anxiety in PTSD participants

The comparison between the severity of PTSD, depression and anxiety in the PTSD group is presented in Figure 1. The total sum of points from BDI-II is originally assembled in four subscores in the following order: minimal depression (0–11 points); mild depression (12–19 points); moderate depression (20–26 points); and severe depression ( $\geq 27$  points). However, since HADS gives three subscores: (1) no symptoms (0–7 points); (2) moderate symptoms (8–10 points); and (3) severe symptoms (11–21 points), for the purpose of this study the first two subscores from BDI-II have been summarized into one combined subscore, to ensure data comparability. The PTSD severity in participants, evaluated by CGI-S, ranged between the following four subscores: mildly ill, moderately ill, markedly ill and severely ill. For the purpose of this study, the last two subscores (markedly ill and severely ill) have been summarized into one combined subscore. Statistical tests confirmed a strong positive correlation between the anxiety (HADS-A) and depression severity index (HADS-D) obtained by HADS alone ( $r = 0.741$ ,  $P < 0.05$ ). Positive correlations were also observed between the results obtained by the three different scales: BDI-II versus HADS-A ( $r = 0.835$ ,  $P < 0.05$ ), BDI-II versus HADS-D ( $r = 0.716$ ,  $P < 0.05$ ) and between CGI-S versus BDI-II ( $r = 0.53$ ,  $P < 0.05$ ).

### Antioxidant enzymes’ activity and MDA concentrations in the PTSD and control group

Data for the eSOD and eGPx catalytic concentrations and sMDA concentration of the control and the PTSD group are shown in Figure 2. Values for catalytic concentrations of antioxidative enzymes are expressed as U/mg Hb in order to avoid possible errors resulting from erythrocyte washing and dilution. Mass concentrations of Hb were within reference values and did not differ significantly between the two groups. A significant difference was noted between PTSD and the control group in eSOD ( $t = 4.1307$ ;  $P = 0.000090$ ) and eGPx ( $t = 3.53797$ ;  $P = 0.000683$ ) activities, with lower values exhibited in the PTSD group. No significant differences were found in sMDA concentrations. The data analysis was further



**Table 1.** Sociodemographic characteristics of the study participants.

Variables	Control (n = 30)	PTSD (n = 50)	Statistics
Age (years) (mean $\pm$ SD)	46.2 $\pm$ 6.3	47.1 $\pm$ 6.4	<sup>a</sup> t = -0.643; P = 0.522
Education level (%)			<sup>b</sup> $\chi^2$ = 28.982; P < 0.0001
$\leq$ 8 years of school	6.7	10.2	
9–12 years of school	26.7	81.6	
> 12 years of school	66.6	8.2	
Employment status (%)			<sup>b</sup> $\chi^2$ = 43.773; P < 0.0001
Full time	86.7	12.2	
Unemployed	0	32.7	
Retired	13.3	55.1	
Exposure time (%)			<sup>b</sup> $\chi^2$ = 2.275; P = 0.3206
1 to 6 months	13.8	6.1	
7–12 months	17.2	20.4	
1–2 years	17.2	28.6	
More than 2 years	48.3	44.9	
Time since exposure (years)	15 $\pm$ 2	15 $\pm$ 2	
Oil consumption (%)			<sup>b</sup> $\chi^2$ = 8.864; P = 0.0119
Sunflower	46	69*	
Olive	28	4*	
Others	26	27	
Fish consumption (%)			<sup>b</sup> $\chi^2$ = 1.043; P = 0.5936
Rarely	36.7	40.8	
Regularly	60.0	51.0	
Often	3.3	8.2	
BMI (kg/m <sup>2</sup> ) (mean $\pm$ SD)	27.0 $\pm$ 4.7	27.3 $\pm$ 3.9	<sup>a</sup> t = -0.001; P = 0.999
TCH (mmol/L) (mean $\pm$ SD)	5.67 $\pm$ 1.23	5.45 $\pm$ 1.14	<sup>a</sup> t = 0.787; P = 0.6559
LDL-CH (mmol/L) (mean $\pm$ SD)	3.37 $\pm$ 0.77	3.42 $\pm$ 0.87	<sup>a</sup> t = -0.218; P = 0.5117
HDL-CH (mmol/L) (mean $\pm$ SD)	1.26 $\pm$ 0.27	1.24 $\pm$ 0.32	<sup>a</sup> t = 0.249; P = 0.3233
TG (mmol/L) (mean $\pm$ SD)	1.95 $\pm$ 0.89	1.87 $\pm$ 1.09	<sup>a</sup> t = 0.349; P = 0.2471

BMI: body mass index; TCH: total cholesterol; LDL-CH: low-density lipoprotein cholesterol; HDL-CH: high-density lipoprotein cholesterol; TG: triglycerides.

<sup>a</sup>Student's *t*-test.

<sup>b</sup>Chi-square test.

\*P < 0.05


focused on relations between the component traits of (anti)oxidative status and serum lipids concentration. A statistically significant correlation was only found for eSOD activity versus serum TCH concentration ( $r = -0.3322$ ,  $P < 0.0500$ ) for patients with PTSD. This relationship was not confirmed neither for the control group ( $r = 0.1384$ ;  $P > 0.0500$ ), nor for the total group ( $r = 0.0161$ ;  $P > 0.0500$ ). Other tested correlations were also not significant (all  $P > 0.005$ , data not shown).

## Discussion


Very few studies have been conducted on patients afflicted with a combat-related PTSD, and to the best of our knowledge, none of them analysed the

relationship between parameters of oxidative stress and the severity of clinical symptoms.

Fifty patients who met the DSM-IV criteria for PTSD but did not have any previously documented somatic illnesses were recruited among outpatients in the University Psychiatric Hospital Vrapče, Zagreb, Croatia. Traumatic experiences during the Homeland War acted as a trigger for PTSD development for all of them. The control group was matched with 30 healthy volunteers who also served in the Homeland War, fortunately without serious consequences on their mental health. All participants were males of similar age, and the time spent in the war did not differ significantly between groups. Thus, two homogeneous groups were obtained according to gender, age, stressor and duration of illness.



**Figure 1.** Comparison between the scores in PTSD patients obtained by Beck Depression Inventory (BDI-II), Hospital Anxiety and Depression Scale for anxiety (HADS-A) and depression (HADS-D), and Clinical Global Impression-Severity scale (CGI-S). Values are expressed as percentage of total group.



**Figure 2.** Erythrocyte superoxide dismutase (eSOD) and glutathione peroxidase (eGPx) catalytic concentrations and serum lipid peroxide (sMDA) concentration in patients with PTSD and controls included in the study. Values are expressed as mean  $\pm$  SD. \* $P < 0.05$ .

The adverse effects and the intensity of the damage caused by free radicals strongly depend on the protective action of enzymatic (i.e. SOD, GPx and CAT) as well as non-enzymatic antioxidants (i.e. vitamins A, C and E, glutathione and selenium). In our study, the catalytic concentrations of eSOD and eGPx were

within the reference range for both groups, but statistically significant differences for eSOD and eGPx activity between the two groups were found, with lower values obtained for the PTSD group. Similar results for SOD and GPx were obtained by Dadheech *et al.*<sup>41</sup> in blood of patients with schizophrenia, while

Akyol *et al.*<sup>42</sup> reported decreased SOD with unchanged GPx catalytic concentrations.

Superoxide dismutase and GPx are related enzymes since the latter removes hydrogen peroxide, produced by SOD during superoxide radical dismutation. A positive correlation between SOD and GPx was demonstrated when tested in all participants as a whole ( $r=0.56$ ;  $P<0.05$ ). This positive correlation between SOD and GPx was again confirmed when tested within each study group separately ( $r=0.50$ ;  $P<0.05$  for PTSD group, and  $r=0.43$ ;  $P<0.05$  for control group).

These results could be explained through the data described by Zieker *et al.*<sup>43</sup> who referred different gene expression pattern in patients with PTSD, with SOD being downregulated. In addition, genes for some interleukins and thioredoxin, another protein involved in antioxidative defence, were also downregulated. In contrast to the above mentioned results, Tezcan *et al.*<sup>44</sup> did not find any difference in SOD and GPx activities between patients with PTSD and healthy controls. In literature, there are reports on lower SOD activity being related to aging processes and some other disorders.<sup>45,46</sup> Attention is drawn to the negative correlation between eSOD activity and cholesterol concentration, specifically in the PTSD group. Although similar relationships were observed in human and animal studies with nutritional interventions with different types and amounts of fats,<sup>47,48</sup> available data are ambiguous and the final explanation of related mechanisms is yet to be found.

Oxidative stress is demonstrated by increased concentrations of plasma peroxides.<sup>49</sup> Dadheech *et al.*<sup>41</sup> and Akyol *et al.*<sup>42</sup> have found increased oxidative stress in patients with schizophrenia as indicated by higher MDA concentrations. However, in our study, concentration of sMDA did not differ significantly between the two groups. Likewise, Tezcan *et al.*<sup>44</sup> did not find any increase in MDA concentration in the serum of PTSD patients.

The principal targets of lipid peroxidation are unsaturated fatty acids originating mainly from food. Therefore, some nutritional habits of the participants were compared, i.e. oil type and frequency of fish consumption. Generally, no significant differences were found between the two groups, with the exception of oil type consumption. Although the control group consumed more olive oil, no correlations with serum lipid parameters, antioxidant enzyme activities and sMDA concentrations were found. Additionally, age, time spent in the war, BMI and fish consumption did not correlate significantly with parameters of oxidative stress ( $P>0.05$  for all correlations) (data not shown).

Since anxiety and depression are the most common co-morbid symptoms in PTSD, self-report

questionnaires were given to the affected participants. More than 50% of them reported severe and about 20% reported moderate symptoms of depression in BDI-II as well as in HADS. Separate scores for depression and anxiety from HADS revealed anxiety symptoms to be more pronounced. Most of the participants with PTSD were moderately ill when evaluated by the clinician-rated scale. Observed discrepancies between clinician-rated and self-administered severity of illness were reported in a number of studies.<sup>50</sup>

It is worth noting that the education level was significantly higher in the control compared to the PTSD group, and higher education may indicate better cognitive abilities. Similarly, Horner and Hamner<sup>51</sup> in their review of performance on standard neuropsychological tests among individuals with PTSD suggested that certain individuals who are at risk for developing PTSD are also characterized by subtle cognitive deficits. Whether or not, and how, the level of education and cognitive abilities can be related to the likelihood that certain trauma will result in PTSD development, still remains to be investigated. Furthermore, because of their disease, the majority of PTSD patients lost their jobs, and/or were forced to retire. Lower incomes undoubtedly affected their social status as well as lifestyle and nutritional habits which resulted in lower nutritional quality, as shown by oil type consumption, although the higher BMI suggests that the quantity of food intake was not lowered. As reported by Braš *et al.*,<sup>52</sup> PTSD symptoms were associated with a lower quality of life in the affected population. Since it had not been possible to analyse the quality of nutrition at the moment of the disease onset no conclusions on this topic can be attempted, but bearing in mind the impact of certain nutrients like vitamins, selenium, iron and others on (anti)oxidative status, these relationships certainly deserve further attention.

Finally, some drawbacks of this study must be pointed out. The main one is the fact that the PTSD patients were on antidepressant therapy with sertraline, which might cause, through the alleviation of depressive symptoms, a reduction of plasma markers of oxidative stress.<sup>53</sup> However, the patients could have risked further exacerbation of clinical symptoms had this antidepressant been removed from their therapy. In order to avoid the dose effect, all the patients were on the same dose (50 mg/day). In addition, changes in the regulation of the HPA axis could be followed. Gotovac *et al.*<sup>54</sup> have shown in their longitudinal study on a similar group of PTSD patients initially increased cortisol concentrations and natural killer cell cytotoxicity. However, cortisol concentrations decreased with the passing of time after a traumatic experience. It is worth noting that the changes in cortisol concentration reported in the above mentioned study were still within the physiological



range, as were the reported significantly lower activities of sSOD and sGPx, although still within reference values. Furthermore, only a few nutritional questions were included. Further, more detailed studies about the influence of nutritional and environmental factors on parameters of oxidative stress should bring new insights into mechanisms of the disease onset and development.

In conclusion, the presence of increased oxidative stress indicators in patients suffering from PTSD could not be confirmed. However, our data suggest that lower catalytic concentrations of eSOD and eGPx in patients with PTSD may indicate a weakened response to oxidative stress due to impaired enzyme activity and/or decreased enzyme synthesis. Although no changes in MDA concentrations were found at the time of this study, presumably oxidative stress was present soon after the traumatic experience. Our results suggest a compensated balance between lipid peroxidation and enzymatic activity, though the lower level of oxidative stress due to therapy with antidepressants (sertraline) must also be taken into account. Finally, oxidative stress might be involved in the onset of PTSD, but is not essentially responsible for the continuance of the illness. Further studies are necessary to reveal the underlying mechanisms in PTSD development and possible treatments.

### Acknowledgements

The authors thank all the volunteers who participated in the study. The authors also wish to thank Mr. Goran Kapustić and Mr. Franjo Ljubić for their skilful assistance, and Mrs. Laurette Rako and Ms. Dijana Gugić for manuscript editing and language advice.

### Declaration of conflicting interests

None declared.

### Funding

This research was funded by the grant from the Croatian Academy of Sciences and Arts.

### Ethical approval

The study protocol was approved by the Ethics Committees of the University Psychiatric Hospital Vrapče and the School of Medicine, University of Zagreb, Croatia.

### Guarantor

ID.

### Contributorship

LBŠ, DK and ID researched literature and conceived the study. LBŠ, BBLj, MMP and MČ performed laboratory

analyses. LBŠ, DK, ID and NM were involved in protocol development, gaining ethical approval, patient recruitment and interpretation data analysis. JA and ID undertook statistical analysis. LBŠ wrote the first draft of the manuscript. PFG was involved in patient treatment and recruitment. All authors reviewed and edited the manuscript and approved the final version of the manuscript. LBŠ and DK have contributed equally to this work.

### References

1. Abuja PM and Albertini R. Methods for monitoring oxidative stress, lipid peroxidation and oxidation resistance of lipoproteins. *Clin Chim Acta* 2001; 306: 1–17.
2. Benzie IF. Lipid peroxidation: a review of causes, consequences, measurement and dietary influences. *Int J Food Sci Nutr* 1996; 47: 233–261.
3. Powers SK, De Ruisseau KC, Quindry J, et al. Dietary antioxidants and exercise. *J Sports Sci* 2004; 22: 81–94.
4. Halliwell B. Oxidative stress and neurodegeneration: where are we now? *J Neurochem* 2006; 97: 1634–1658.
5. Mahadik SP, Evans D and Lal H. Oxidative stress and role of antioxidant and  $\omega$ -3 essential fatty acid supplementation in schizophrenia. *Prog Neuropsychopharmacol Biol Psychiat* 2001; 25: 463–493.
6. Mahadik SP, Pilai A, Joshi S, et al. Prevention of oxidative stress-mediated neuropathology and improved clinical outcome by adjunctive use of a combination of antioxidants and omega-3 fatty acids in schizophrenia. *Int Rev Psych* 2006; 18: 119–131.
7. Tylec A, Jarzab A, Stryjecka-Zimmer M, et al. Stress oxidative in schizophrenia. *Pol Merkuriusz Lek* 2007; 23: 74–77.
8. Valko M, Leibfritz D, Moncol J, et al. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007; 39: 44–84.
9. Lebel C. Oxygen radicals: common mediators of neurotoxicity. *Neurotox Teratol* 1991; 13: 341–346.
10. Ranjekar PK, Hinge A, Hegde MV, et al. Decreased antioxidant enzymes and membrane essential polyunsaturated fatty acids in schizophrenic and bipolar mood disorders patients. *Psych Res* 2003; 121: 109–122.
11. Freeman MP. Omega 3 fatty acids in psychiatry: a review. *Ann Clin Psych* 2000; 12: 159–165.
12. Mandelsamen Perica M and Delaš I. Essential fatty acids and psychiatric disorders. *Nutr Clin Pract* 2011; 26: 409–425.
13. Kalinić D, Borovac Štefanović L, Jerončić A, et al. Eicosapentaenoic acid in serum lipids could be inversely correlated with severity of clinical symptomatology in Croatian war veterans with posttraumatic stress disorder. *Croat Med J* 2014; 55: 27–37.
14. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, Fifth Edition. Arlington, VA: American Psychiatric Association, 2013. Available from [dsm.psychiatryonline.org](http://dsm.psychiatryonline.org) (accessed 13 March 2014).
15. Evans DR, Parikh VV, Khan MM, et al. Red blood cell membrane essential fatty acid metabolism in early psychotic patients following antipsychotic drug treatment.

- Prostaglandins Leukot Essent Fatty Acids* 2003; 69: 393–399.
16. Skinner AO, Mahadik SP and Garver DL. Thiobarbituric acid reactive substances in the cerebrospinal fluid in schizophrenia. *Schizophr Res* 2005; 76: 83–87.
  17. Chakraborty S, Singh OP, Dasgupta A, et al. Correlation between lipid peroxidation-induced TBARS level and disease severity in obsessive-compulsive disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33: 363–366.
  18. Kuloglu M, Ustundag B, Atmaca M, et al. Lipid peroxidation and antioxidant enzyme levels in patients with schizophrenia and bipolar disorder. *Cell Biochem Funct* 2002; 20: 171–175.
  19. Hui-Chun L, Qiao-Zhen C, Ying MA, et al. Imbalanced free radicals and antioxidant defense systems in schizophrenia: a comparative study. *J Zhejiang Univ Sci B* 2006; 7: 981–986.
  20. Hovatta I, Juhila J and Donner J. Oxidative stress in anxiety and comorbid disorders. *Neurosci Res* 2010; 68: 261–275.
  21. Sala M, Perez J, Soloff P, et al. Stress and hippocampal abnormalities in psychiatric disorders. *Eur Neuropsychopharmacol* 2004; 14: 393–405.
  22. Sternberg EM. Neural regulation of innate immunity: a coordinated nonspecific host response to pathogens. *Nat Rev Immunol* 2006; 6: 318–328.
  23. Kuloglu M, Atmaca M, Tezcan E, et al. Antioxidant enzyme activities and malondialdehyde levels in patients with obsessive-compulsive disorder. *Neuropsychobiology* 2002; 46: 27–32.
  24. Kuloglu M, Atmaca M, Tezcan E, et al. Antioxidant enzyme and malondialdehyde levels in patients with panic disorder. *Neuropsychobiology* 2002; 46: 186–189.
  25. Bouayed J, Rammal H and Soulimani R. Oxidative stress and anxiety. Relationship and cellular pathways. *Oxid Med Cell Longev* 2009; 2: 63–67.
  26. Falkai P and Möller H-J. Psychopathology: genetics and the stress-vulnerability hypothesis. *Eur Arch Psychiatry Clin Neurosci* 2012; 262: 181–182.
  27. Švob Štrac D, Mustapić M, Šagud M, et al. Lipid levels in neuropsychiatric disorders. In: Araujo C and Perez D (eds) *Triglycerides: chemical structure, biosynthesis and role in disease*. New York: Nova Science Publishers, Inc., 2013, pp.1–79.
  28. Linden SC, Hess V and Jones E. The neurological manifestations of trauma: lessons from World War I. *Eur Arch Psychiatry Clin Neurosci* 2012; 262: 253–264.
  29. American Psychiatric Association (APA) (Ed.). *Diagnostic and statistical manual of mental disorders (DSM-IV)*, 4th ed., text rev. Washington, DC: American Psychiatric Association Press, 2000.
  30. WHO. *International Statistical Classification of Diseases and Related Health Problems ICD-10 (Meunarodna klasifikacija bolesti i srodnih zdravstvenih problema – MKB-10, in Croatian)*. 10th revision, Vol 1, 2nd ed. Zagreb, Croatia: Medicinska naklada, 2012.
  31. Zigmond AS and Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983; 67: 361–370.
  32. Beck AT, Steer RA, Ball R, et al. Comparison of beck depression inventories -IA and -II in psychiatric outpatients. *J Pers Assess* 1996; 67: 588–597.
  33. National Institute of Mental Health. CGI Clinical Global Impressions. In: Guy W (ed.) *Early Clinical Drug Evaluation Unit (ECDEU) assessment manual for psychopharmacology, Revised*. (NIMH publication DHEW publ. No (ADM) 76-338) Bethesda, MD: National Institute of Mental Health, 1976, pp.218–222.
  34. Senta A, Pucarín-Cvetković J and Doko Jelinić J. *Quantitative models of food and meals, (in Croatian)*. Zagreb, Croatia: Medicinska naklada, 2004.
  35. Paglia DE and Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 1967; 70: 158–169.
  36. Woolliams JA, Wiener G, Anderson PH, et al. Variation in the activities of glutathione-peroxidase and superoxide-dismutase and in the concentration of copper in the blood in various breed crosses of sheep. *Res Vet Sci* 1983; 34: 253–256.
  37. Anguloa J, Romera JM, Ramirez M, et al. Effects of storage conditions on lipid oxidation in infant formulas based on several protein sources. *JAOCS* 1998; 75: 1603–1607.
  38. Botsoglou NA, Fletouris DJ, Papageorgiou GE, et al. Rapid, sensitive and specific thiobarbituric acid method for measuring lipid peroxidation in animal tissue, food and feedstuff samples. *J Agric Food Chem* 1994; 42: 1931–1937.
  39. Xie W-L and Ji J-M. Antioxidant activities of vitamins E and C in a novel liposome system. *J Food Biochem* 2008; 32: 766–781.
  40. Friedewald WT, Levy RI and Fredrickson DS. Estimation of the concentration of low density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; 18: 499–502.
  41. Dadheech G, Mishra S, Gautam S, et al. Evaluation of antioxidant deficit in schizophrenia. *Indian J Psychiatry* 2008; 50: 16–20.
  42. Akyol O, Herken H, Uz E, et al. The indices of endogenous oxidative and antioxidative processes in plasma from schizophrenic patients. The possible role of oxidant/antioxidant imbalance. *Prog Neuropsychopharmacol Biol Psychiatry* 2002; 26: 995–1005.
  43. Zieker J, Zieker D, Jatzko A, et al. Differential gene expression in peripheral blood of patients suffering from post-traumatic stress disorder. *Mol Psychiatry* 2007; 12: 116–118.
  44. Tezcan E, Atmaca M, Kuloglu M, et al. Free radicals in patients with post-traumatic stress disorder. *Eur Arch Psychiatry Clin Neurosci* 2003; 253: 89–91.
  45. Kumawat M, Sharma TK, Singh I, et al. Decrease in antioxidant status of plasma and erythrocytes from geriatric population. *Dis Markers* 2012; 33: 303–308.
  46. Laškaj R, Dodig S, Čepelak I, et al. Superoxide dismutase, copper and zinc concentrations in platelet-rich plasma of pneumonia patients. *Ann Clin Biochem* 2009; 46: 123–128.

47. Yuan YV, Kitts DD and Godin DV. Variations in dietary fat and cholesterol intakes modify antioxidant status of SHR and WKY Rats. *J Nutr* 1998; 128: 1620–1630.
48. Mohammadi M, Alipour M, Alipour MR, et al. Effects of high cholesterol diet and parallel chronic exercise on erythrocyte primary antioxidant enzymes and plasma total antioxidant capacity in Dutch rabbits. *Int J Endocrinol Metab* 2006; 4: 30–40.
49. Maes M, Galecki P, Chang YS, et al. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; 35: 676–692.
50. Enns MW, Larsen DH and Cox BJ. Discrepancies between self and observer rating of depression. The relationship to demographic, clinical and personality variables. *J Affect Dis* 2000; 60: 33–41.
51. Horner MD and Hamner MB. Neurocognitive functioning in posttraumatic stress disorder. *Neuropsychol Rev* 2002; 12: 15–30.
52. Braš M, Milunović V, Boban M, et al. A quality of life in chronic combat related PTSD – A study on Croatian war veterans. *Coll Antropol* 2011; 35: 681–686.
53. Michalakeas CA, Parissis JT, Douzenis A, et al. Effects of sertraline on circulating markers of oxidative stress in depressed patients with chronic heart failure: a pilot study. *J Card Fail* 2011; 17: 748–754.
54. Gotovac K, Sabioncello A, Rabatic S, et al. Flow cytometric determination of glucocorticoid receptor (GCR) expression in lymphocyte subpopulations: lower quantity of GCR in patients with post-traumatic stress disorder (PTSD). *Clin Exp Immunol* 2003; 131: 335–339.