

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 25, 2015, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series historia naturalis, 25, 2015, 2

ISSN 1408-533X

UDK 5

Letnik 25, leto 2015, številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Dunja Bandelj Mavsar, Nicola Bettoso (IT), Christian Capapé (F), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Mitja Kaligarič, Gregor Kovačič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Alenka Malej, Patricija Mozetič, Martina Orlando - Bonaca, Michael Stachowitsch (A), Tom Turk, Elena Varljen Bužan

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Patricija Mozetič

Lektor/Supervisione/Language editor:

Polona Šergon (sl.), Petra Berlot (angl.)

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Prelom/Composizione/Typesetting:

Grafis trade d.o.o.

Tisk/Stampa/Print:

Grafis trade d.o.o.

Izdajatelj/Editore/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria©

**Za izdajatelja/Per Editore/
Publisher represented by:**

Salvator Žitko

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 20. 12. 2015.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS)

Annales - series historia naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation:

300 izvodov/copie/copies

Revija Annales series historia naturalis je vključena v naslednje podatkovne baze: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL).

VSEBINA / INDICE GENERALE / CONTENTS

RECENTNE SPREMEMBE
V SREDOZEMSKI IHTIOFAVNI
CAMBIAMENTI RECENTI
NELLA ITTIOFAUNA MEDITERRANEA
RECENT CHANGES
IN THE MEDITERRANEAN ICHTHYOFAUNA

Sihem RAFRAFI-NOUIRA, Olfa EL KAMEL-MOUTALIBI, Mohamed Mourad BEN AMOR, Christian CAPAPÉ
Additional records of Spinetail devilray *Mobula japanica* (Chondrichthyes: Mobulidae) from the Tunisian coast (Central Mediterranean) 103
Nov zapis o pojavljanju mante vrste *Mobula japanica* (Chondrichthyes: Mobulidae) vzdolž tunizijske obale (osrednji Sredozemlje)

Okan AKYOL & İlker AYDİN
Additional records of two lesepsian fish, *Siganus luridus* and *Champsodon vorax* from Izmir Bay (Aegean sea, Turkey) 109
Dodatni zapisi o pojavljanju dveh vrst lesepskih ribjih selivk, *Siganus luridus* in *Champsodon vorax*, iz Izmirskega zaliva (Egejsko morje, Turčija)

SREDOZEMSKI MORSKI PSI
SQUALI DEL MEDITERRANEO
MEDITERRANEAN SHARKS

Hakan KABASAKAL & Sait Özgür GEDİKOĞLU
Shark attacks against humans and boats in Turkey's waters in the twentieth century 115
Napadi morskih psov na ljudi in plovila v turških vodah v dvajsetem stoletju

Halit FİLİZ & Hakan KABASAKAL
Photographic record of the Spinner shark, *Carcharhinus brevipinna* (Müller & Henle, 1839), in Gokova Bay (south Aegean Sea, Turkey) 123
Fotografski zapis o kratkoplavutem morskem psu, *Carcharhinus brevipinna* (Müller & Henle, 1839), v zalivu Gokova Bay (južno Egejsko Morje, Turčija)

IHTIOLOGIJA
ITTIOLOGIA
ICHTHYOLOGY

Balkis SALLAMI, Mohamed BEN SALEM, Sihem RAFRAFI-NOUIRA, Olfa EL KAMEL-MOUTALIBI, Christian REYNAUD & Christian CAPAPÉ
Observations on Thinlip Conger *Gnathophis mystax* (Osteichthyes: Congridae) from the Tunisian Coast (Central Mediterranean) 131
Opazovanja dolgonosega ugorja *Gnathophis mystax* (Osteichthyes: Congridae) ob tunizijski obali (osrednji Mediteran)

Nicola BETTOSO & Giovanni COMISSO
Recent record of the Serpent Eel *Ophisurus serpens* (Ophichthidae) in the Gulf of Trieste (Northern Adriatic Sea) 141
Nov zapis o pojavljanju zobate jegulje *Ophisurus serpens* (Pisces: Ophichthidae) v Tržaškem zalivu (Severni Jadran)

Jakov DULČIĆ & Pero TUTMAN
Additional record of Common bream *Abramis brama* (Cyprinidae) in the Adriatic drainage system (Norin River, Croatia) 145
Novi podatek o pojavljanju ploščiča (*Abramis brama*, Cyprinidae) v Jadranskem povodju (reka Norin, Hrvaška)

FAVNA
FAUNA
FAUNA

Toni KOREN & Domen TRKOV
Contribution to the Scarabeoidea (Coleoptera) of island Cres, Croatia 151
Prispevek k poznavanju favne Scarabaeoidea (Coleoptera) otoka Cres, Hrvaška

MISCELLANEA

**Dejan PALISKA, Simon KERMA,
Rudi ČOP & Flavio BONIN**

An attempt to demonstrate the influence
of Maunder Minimum Climate on salt
production and its price in the Slovenian Istria
(Sečovlje Salt-Pans) 163
*Poskus prikaza vpliva podnebja iz obdobja
Maunderjevega minimuma na
proizvodnjo soli in njeno ceno
v Slovenski Istri (Sečoveljske soline)*

Danijel IVAJNŠIČ & Mitja KALIGARIČ

Can evapotranspiration be considered
an additional indicator for understanding
the changed landscape identity
of the classic Karst? 173
*Ali lahko evapotranspiracijo smatramo
kot dodatni pokazatelj za razumevanje
spremenjene identitete klasičnega Krasa?*

DELO NAŠIH ZAVODOV IN DRUŠTEV
*ATTIVITÀ DEI NOSTRI ISTITUTI
E DELLE NOSTRE SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS
AND ASSOCIATIONS*

22. Simpozij o okoljski biogeokemiji v Piranu
(**Jadran Faganeli in Nives Ogrinc**) 185

OCENE IN POROČILA
*RECENSIONI E RELAZIONI
REVIEWS AND REPORTS*

Ocena knjige: »ENDEMI U HRVATSKOJ FLORI«
avtorjev Tonija Nikolića, Milenka Milovića,
Sandra Bogdanovića in Nenada Jasprice
(**Mitja Kaligarič**) 189

Anton Brancelj: JAMA VELIKA PASICA:
ZGODOVINA, OKOLJE IN ŽIVLJENJE V NJEJ
/ *THE VELIKA PASICA CAVE: THE HISTORY,
ENVIRONMENT AND LIFE IN IT*. Založba ZRC
in Nacionalni inštitut za biologijo, Ljubljana,
2015, 110 str. (**Lovrenc Lipej**) 190

Navodila avtorjem 193
Istruzioni per gli autori 195
Instructions to authors 197

Kazalo k slikam na ovitku 200
Index to images on the cover 200

Short scientific article
Received: 2015-12-01

UDC 597.551.2:591.9(282.249.1)

ADDITIONAL RECORD OF COMMON BREAM *ABRAMIS BRAMA* (CYPRINIDAE) IN THE ADRIATIC DRAINAGE SYSTEM (NORIN RIVER, CROATIA)

Jakov DULČIĆ & Pero TUTMAN

Institute of Oceanography and Fisheries, POB 500, HR-21000 Split, Croatia
E-mail: dulcic@izor.hr

ABSTRACT

On 14 March 2015 an adult specimen of common bream Abramis brama was caught with fish trap in Norin River (right bank tributary of the Neretva River, Adriatic drainage system, near settlement Vid, Croatia). This represents the second record of this species for the Adriatic drainage system in Croatia.

Key words: Cyprinidae, *Abramis brama*, Adriatic drainage system, Croatia

NUOVE SEGNALAZIONI DELL'ABRAMIDE COMUNE *ABRAMIS BRAMA* (CYPRINIDAE) NEL SISTEMA DI DRENAGGIO ADRIATICO (FIUME NORIN, CROAZIA)

SINTESI

Il 14 marzo 2015 un esemplare adulto dell'abramide comune (Abramis brama) è stato catturato con una nassa nel fiume Norin (affluente della riva destra del fiume Neretva, sistema di drenaggio dell'Adriatico, vicino al villaggio di Vid, in Croazia). Questa cattura rappresenta la seconda segnalazione della specie nel sistema di drenaggio adriatico in Croazia.

Parole chiave: Cyprinidae, *Abramis brama*, sistema di drenaggio dell'Adriatico, Croazia

INTRODUCTION

The common bream, *Abramis brama* (Linnaeus, 1758) (Cyprinidae), inhabits most European drainages from Adour (France) to Pechora (White Sea basin); Aegean Sea basin, in Lake Volvi and Struma and Maritza drainages (Kottelat & Freyhof, 2007). It is not native to Iberian Peninsula, Adriatic basin, Italy, Scotland, and Scandinavia north of Bergen (Norway) and 67°N (Finland). It is locally introduced in Ireland, Spain, north-eastern Italy, from Marmara basin (Turkey) and eastward to Aral basin, in Lake Baikal and upper Ob and Yenisei drainages (Kottelat & Freyhof, 2007). In Croatia, this species is found only in the waters of the Black Sea drainage system (Glamuzina *et al.*, 2013).

MATERIAL AND METHODS

On 14 March 2015 an adult specimen (Fig. 1) of common bream was caught with fish trap in Norin River (right bank tributary of the Neretva River, Adriatic drainage system, near settlement Vid, Croatia) (43.081644°N, 17.629486°E). This represents the second record of this species for the Adriatic drainage system in Croatia.

RESULTS AND DISCUSSION

The first record of the common bream for the Adriatic drainage system (Mala Neretva River, wider area of Neretva River estuary, Croatia) was on 17 April 2010 (male, total length = 43.8 cm, weight = 1047 g) (Bartulović *et al.*, 2010).

One question could arise after this additional record in the wider area of Neretva River estuary: "Has this species established a population?" Although there is still no

evidence of a permanent population in the study area (not enough available reports confirmed on a scientific basis), the capture described here might be an indication of that since some fishermen have signalled the species in some parts of Neretva river delta during 2015 (Dugandžić, *pers. comm.*). The only possible explanation for such record is a not sufficiently controlled introduction (in this case of a non-native species).

The common bream has been introduced to the Neretva River and now represents a potential threat to the natural equilibrium of their community. This species may develop stunted high density populations becoming locally abundant, with potential negative consequences both within and beyond the local fish community due to competition for food resources or hybridization (Bartulović *et al.*, 2010). Furthermore, common bream often has a pronounced migratory behaviour and may consequently perform considerable distances to lakes within a river system (Volta *et al.*, 2013). Successful non-native species are often characterized by high physiological tolerance and functional characteristics different from those of the members of invaded communities and have been reported to affect the functional diversity of communities with possible strong impacts on food webs and ecosystem functioning (see Bartulović *et al.*, 2010). These features make the common bream a potentially effective and highly undesirable invader of southern European waters (see Volta *et al.*, 2013).

It is quite clear that non-native species can have significant effects on the composition of entire communities by displacement of local species with similar trophic level, by altering the behaviour or habitat selection of prey, resulting in a significant disturbance of the local communities interactions (Cucherousset & Olden, 2011). However, at this point it is not known to what



Fig. 1: (Left) *Abramis brama* caught in Norin River (Croatian coast, Adriatic drainage system); (right) male, with nuptial tubercles on the head. (Photo: B. Markota)

Sl. 1: Ploščič (*Abramis brama*), ujet v reki Norin (hrvaška obala, jadransko povodje) (levo); samec, dobro vidni paritveni grebenčki na glavi (desno). (Foto: B. Markota)

extent the occurrence of the common bream in the Neretva watershed is related to environmental quality and to natural biodiversity. Therefore it is evident that understanding on all aspects of the invasion process, from introduction to the establishment, spread and impacts is still required.

Prevention measures should be taken by the administration to avoid the spreading of this species to other reservoirs and river basins. Eradication of introduced fish

is practically impossible in large freshwater ecosystems, so prevention of further introductions and translocations is of primary concern.

ACKNOWLEDGEMENT

We thank Mr. Branko Markota (Vid, Croatia) for providing photos and data on specimen of the common bream.

NOVI PODATEK O POJAVLJANJU PLOŠČIČA (*ABRAMIS BRAMA*, CYPRINIDAE) V JADRANSKEM POVODJU (REKA NORIN, HRVAŠKA)

Jakov DULČIĆ & Pero TUTMAN

Institute of Oceanography and Fisheries, POB 500, HR-21000 Split, Croatia

E-mail: dulcic@izor.hr

POVZETEK

14. marca 2015 je bil v ribiško mrežo ujet odrasel primerek ploščiča (*Abramis brama*) v reki Norin (jadransko povodje, desni pritok reke Neretve pri naselju Vid, Hrvatska). Gre za drugi primer pojavljanja te vrste v jadranskom povodju na Hrvatskem.

Ključne besede: Cyprinidae, *Abramis brama*, jadransko povodje, Hrvatska

REFERENCES

- Bartulović, V., J. Dulčić, I. Bogut, J. Pavličević, E. Hasković & B. Glamuzina (2010):** First record of the freshwater bream, *Abramis brama* in the river Mala Neretva, Adriatic drainage system of Croatia. *Cybium*, 35 (2), 165-166.
- Cucherousset, J. & J. D. Olden (2011):** Ecological impacts of non-native freshwater fishes. *Fisheries*, 36, 215-230.
- Glamuzina, B., J. Pavličević, P. Tutman, L. Glamuzina, I. Bogut & J. Dulčić (2013):** Ribe Neretve. Udruga CEAV - Centar za zaštitu i promicanje endemskih i autohtonih ribljih vrsta, Mostar, Republika Bosna i Hercegovina, Mostar /Metković, Modrozeleno - Zadruga branitelja, Metković, 261 str.
- Kottelat, M. & J. Freyhof (2007):** Handbook of European Freshwater Fishes. Kottelat, Cornol, Switzerland and Freyhof, Berlin, Germany, 646 p.
- Volta, P., E. Jeppesen, B. Leoni, B. Campi, P. Sala, L. Garibaldi, L. Lauridsen Torben & I. F. Winfield (2013):** Recent invasion by a non-native cyprinid (common bream *Abramis brama*) is followed by major changes in the ecological quality of a shallow lake in southern Europe. *Biol. Invasions*, 15 (9), 2065-2079.