

# Limit theorems for random Lasota-Yorke maps using the spectral method

Davor Dragičević, UNSW

(joint work with Gary Froyland, Cecilia González-Tokman and  
Sandro Vaienti)

November 28, 2016

D.D was supported by an Australian Research Council Discovery Project DP150100017 and by the Croatian Science Foundation under the project HRZZ-IP-09-2014-2285

# Piecewise expanding maps

Let  $I = [0, 1]$  denote the unit interval equipped with Borel  $\sigma$ -algebra  $\mathcal{B}$  and a Lebesgue measure  $m$ . We say that  $T: I \rightarrow I$  is a *piecewise expanding map* if there exists a partition

$$0 = x_0 < x_1 < \dots < x_{k-1} < x_k = 1$$

and  $\alpha > 1$  such that:

- 1 restriction  $T|_{(x_{i-1}, x_i)}$  is a  $C^1$  function which can be extended to a  $C^1$  function on  $[x_{i-1}, x_i]$ ;
- 2  $|T'(x)| \geq \alpha > 0$  for  $x \in (x_{i-1}, x_i)$ ;
- 3  $g(x) = \frac{1}{|T'(x)|}$  is a function of bounded variation.

# Deterministic setting

- ① *transfer operator*:  $L: L^1(m) \rightarrow L^1(m)$  defined by

$$Lf(x) = \sum_{y \in T^{-1}(x)} \frac{f(y)}{|T'(y)|};$$

- ②  $L$  is quasicompact of diagonal type on the  $BV$  space and consequently it has a nonnegative fixed point which gives the existence of *acim*;
- ③ under some additional assumptions *acim* is unique (and we denote it by  $\mu$ ) and mixing;
- ④ in this case  $L = \Pi + N$  and we have the exponential decay of correlation result.

# Central limit theorem

Assume that  $\phi: I \rightarrow \mathbb{R}$  bounded observable in  $BV$  such that  $\int_{[0,1]} \phi d\mu = 0$ . For each  $n \in \mathbb{N}$ , let

$$S_n = \sum_{k=0}^{n-1} \phi \circ T^k.$$

Theorem (Rousseau–Egele, 1983)

We have that  $\lim_{n \rightarrow \infty} \int_{[0,1]} \frac{S_n^2}{n} = \sigma^2$ , where

$$\sigma^2 = \int_{[0,1]} \phi^2 d\mu + 2 \sum_{n=1}^{\infty} \int_{[0,1]} \phi(\phi \circ T^n) d\mu < \infty.$$

If  $\sigma^2 > 0$ , then  $\frac{S_n}{\sqrt{n}}$  converges in distribution to  $N(0, \sigma^2)$ .

# Large deviation principle

## Theorem

*If  $\sigma^2 > 0$ , then there exists  $\delta > 0$  and a strictly convex, continuous and nonnegative function  $c: (-\delta, \delta) \rightarrow \mathbb{R}$  which vanishes only at 0 such that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mu(S_n > n\varepsilon) = -c(\varepsilon), \quad \text{for } \varepsilon \in (0, \delta).$$

# Ideas of the proofs

We define

$$L_\theta(g) = L(e^{\theta\phi}g), \quad \text{for } g \in BV \text{ and } \theta \in \mathbb{C}.$$

Since  $\theta \mapsto L_\theta$  is analytic, for  $\theta$  sufficiently close to 0,

$$L_\theta = \omega(\theta)\Pi(\theta) + N(\theta).$$

For CLT ( $d\mu = f \, dm$ ):

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{[0,1]} e^{itS_n/\sqrt{n}} d\mu &= \lim_{n \rightarrow \infty} \int_{[0,1]} L_{it/\sqrt{n}}^n(f) \, dm = \lim_{n \rightarrow \infty} \omega(it/\sqrt{n})^n \\ &= e^{-t^2\sigma^2/2}, \end{aligned}$$

for  $t \in \mathbb{R}$ .

For LDP:

we first show that  $\omega'(0) = 0$  and  $\omega''(0) = \sigma^2$  and then that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \int_{[0,1]} e^{\theta S_n} d\mu = \Lambda(\theta),$$

where  $\Lambda(\theta) = \log \omega(\theta)$ , for  $\theta \in \mathbb{R}$  sufficiently close to 0.

# Random Lasota-Yorke maps

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space and assume that  $\sigma: \Omega \rightarrow \Omega$  is invertible transformation that preserves  $\mathbb{P}$ . Furthermore, assume that  $\mathbb{P}$  is ergodic. We now take the collection  $T_\omega, \omega \in \Omega$  of piecewise expanding maps. For  $\omega \in \Omega$  and  $n \in \mathbb{N}$ , set

$$T_\omega^n = T_{\sigma^{n-1}\omega} \circ \dots \circ T_{\sigma\omega} \circ T_\omega$$

and

$$L_\omega^n = L_{\sigma^{n-1}\omega} \circ \dots \circ L_{\sigma\omega} \circ L_\omega.$$

The associated skew-product transformation  $\tau: \Omega \times I \rightarrow \Omega \times I$  is given by

$$\tau(\omega, x) = (\sigma\omega, T_\omega x).$$



# Existence and uniqueness of ACIM, Buzzi 2000

Under some mild assumption we have that

$$\|L_\omega h\|_{BV} \leq K(\omega) \|h\|_{BV}, \quad \text{for } h \in BV \text{ and } \log K \in L^1(\mathbb{P});$$

and there exists  $N \in \mathbb{N}$  we have

$$\text{var}(L_\omega^N h) \leq \alpha^N(\omega) \text{var}(h) + K^N(\omega) \|h\|_1,$$

for  $h \in BV$  and a.e.  $\omega \in \Omega$  and with

$$\int_{\Omega} \log \alpha^N(\omega) d\mathbb{P}(\omega) < 0.$$

If for each subinterval  $J \subset I$  and for a.e.  $\omega \in \Omega$ , there exists  $n(\omega) \in \mathbb{N}$  such that

$$\text{essinf}_{x \in [0,1]} (L_\omega^n \mathbf{1}_J) > 0, \quad \text{for } n \geq n(\omega),$$

there exists a unique acim (w.r.t.  $\mathbb{P} \times m$ )  $\mu$  for  $\tau$  such that  $\pi_*\mu = \mathbb{P}$ , where  $\pi: \Omega \times I \rightarrow \Omega$  is a projection. We can regard  $\mu$  as a collection of fiber measures  $\mu_\omega$ ,  $\omega \in \Omega$  on  $I$ . Also, one has fiberwise decay of correlation result.

We consider observables  $\phi: \Omega \times I \rightarrow \mathbb{R}$  such that

$$\text{esssup}_{(\omega,x)} |\phi(\omega, x)| < \infty \quad \text{and} \quad \text{esssup}_\omega \text{var}(\phi(\omega, \cdot)) < \infty.$$

Moreover, we assume that

$$\int_{[0,1]} \phi(\omega, \cdot) d\mu_\omega = 0, \quad \omega \in \Omega.$$

We form Birkhoff sums

$$S_n(\omega, x) = \sum_{i=0}^{n-1} (\phi \circ \tau^i)(\omega, x) = \sum_{i=0}^{n-1} \phi(\sigma^i \omega, T_\omega^i x).$$

We are interested in the quenched type of limit theorems i.e. those that give an information about the asymptotic behaviour of Birkhoff sums w.r.t. to  $\mu_\omega$  for "typical"  $\omega$ .

Previous work:

- 1 Kifer, 1998: quenched limit theorems in random environment but not with spectral method;
- 2 Aimino, Nicol and Vaienti, 2014: spectral method but the base space is assumed to be a Bernoulli shift.

Assume that  $(\Omega, \mathcal{F}, \mathbb{P}, \sigma)$  is a probability space where  $\Omega$  is a Borel subset of a separable, complete metric space. Furthermore, let  $B$  be a Banach space and  $\mathcal{L} = L_\omega, \omega \in \Omega$  a family of bounded linear operators on  $B$  such that the map  $\omega \mapsto L_\omega$  is  $\mathbb{P}$ -continuous. Then, for a.e.  $\omega \in \Omega$ , the following limits exist (and are independent on  $\omega$ )

$$\Lambda(\mathcal{L}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|L_\omega^n\|$$

and

$$\kappa(\mathcal{L}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log ic(L_\omega^n),$$

where  $ic(L_\omega^n) = \inf\{r > 0 :$

$L_\omega^n(B_B)$  can be covered with finitely many balls of radius  $r\}$ .

If  $\kappa(\mathcal{L}) < \Lambda(\mathcal{L})$ , then there exists  $1 \leq l \leq \infty$  and a sequence of Lyapunov exponents

$$\Lambda(\mathcal{L}) = \lambda_1 > \lambda_2 > \dots > \lambda_l > \kappa(\mathcal{L}) \quad (\text{if } 1 \leq l < \infty)$$

or

$$\Lambda(\mathcal{L}) = \lambda_1 > \lambda_2 > \dots \quad \text{and} \quad \lim_{n \rightarrow \infty} \lambda_n = \kappa(\mathcal{L}) \quad (\text{if } l = \infty);$$

and for  $\mathbb{P}$ -almost every  $\omega \in \Omega$  there exists a unique splitting (called the *Oseledets splitting*) of  $B$  into closed subspaces

$$B = V(\omega) \oplus \bigoplus_{j=1}^l Y_j(\omega),$$

depending measurably on  $\omega$  and such that:

- ① For each  $1 \leq j \leq l$ ,  $\dim Y_j(\omega)$  is finite-dimensional,  $Y_j$  is equivariant i.e.  $L_\omega Y_j(\omega) = Y_j(\sigma\omega)$  and for every  $y \in Y_j(\omega) \setminus \{0\}$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|L_\omega^n y\| = \lambda_j.$$

- ②  $V$  is equivariant i.e.  $L_\omega V(\omega) \subseteq V(\sigma\omega)$  and for every  $v \in V(\omega)$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|L_\omega^n v\| \leq \kappa(\mathcal{L}).$$

In order to be able to apply MET, we will require that:  $\Omega$  is a Borel subset of a separable, complete metric space and that

the map  $\omega \rightarrow T_\omega$  has a countable range

We also form a twisted cocycle. More precisely, for  $\omega \in \Omega$  and  $\theta \in \mathbb{C}$ , we define

$$L_\omega^\theta(h) = L_\omega(e^{\theta\phi(\omega, \cdot)} h), \quad h \in BV.$$

### Theorem

For  $\theta \in \mathbb{R}$ , we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \int_{[0,1]} e^{\theta S_n(\omega, \cdot)} d\mu_\omega = \Lambda(\theta),$$

for  $\mathbb{P}$ -a.e.  $\omega \in \Omega$  where  $\Lambda(\theta)$  is a top Lyapunov exponent of the cocycle  $L_\omega^\theta$ ,  $\omega \in \Omega$ .

# Regularity of $\Lambda$

In order to establish the corresponding regularity property of  $\Lambda$  we introduce some additional assumptions. We require that:

- 1 norms of  $L_\omega$  are uniformly bounded;
- 2 densities  $v_\omega^0$  are uniformly bounded from below ( $d\mu_\omega = v_\omega^0 dm$ ) away from zero;
- 3 there exists  $D, \lambda > 0$  such that

$$\|L_\omega^n f\|_{BV} \leq D e^{-\lambda n} \|f\|_{BV},$$

for  $f \in BV$ ,  $\int f dm = 0$ ,  $n \in \mathbb{N}$  and a.e.  $\omega$ .

We briefly sketch the argument that shows that  $\theta \mapsto \Lambda(\theta)$  is of  $C^2$  on a neighborhood of 0.



# Regularity of $\Lambda$

Key points:

- 1 we construct the top space as  $v_\omega^0 + \mathcal{W}^\theta(\omega, \cdot)$  where  $\mathcal{W}^\theta$  is a (unique) solution of  $F(\theta, \mathcal{W}) = 0$ , where

$$F(\theta, \mathcal{W}) = \frac{L_{\sigma^{-1}\omega}^\theta(v_{\sigma^{-1}\omega}^0 + \mathcal{W}(\sigma^{-1}\omega, \cdot))}{\int (L_{\sigma^{-1}\omega}^\theta(v_{\sigma^{-1}\omega}^0 + \mathcal{W}(\sigma^{-1}\omega, \cdot))) dm} - \mathcal{W}(\omega, \cdot) - v_\omega^0,$$

where  $\mathcal{W} \in \mathcal{S}$  and

$$\mathcal{S} := \{\mathcal{W}: \Omega \times I \rightarrow \mathbb{C} : \mathcal{W}(\omega, \cdot) \in BV, \text{esssup}_\omega \|\mathcal{W}(\omega, \cdot)\|_{BV} < \infty\}.$$

- 2  $\Lambda(\theta) = \int \log |\int e^{\theta\phi(\omega, \cdot)} (v_\omega^0 + \mathcal{W}^\theta(\omega, \cdot)) dm| d\mathbb{P}(\omega)$ ;
- 3 for  $\theta$  close to 0, the top Oseledets space of the twisted cocycle  $L_\omega^\theta$  is one-dimensional.

Also,  $\Lambda'(0) = 0$  and  $\Lambda''(0) = \Sigma^2$ , where  $\Sigma^2$  is a variance.

### Theorem (Large deviation principle)

Assume that  $\Sigma^2 > 0$ . Then, there exists  $\varepsilon_0 > 0$  and a function  $c: (-\varepsilon_0, \varepsilon_0) \rightarrow \mathbb{R}$  which is nonnegative, continuous, strictly convex, vanishing only at 0 and such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mu_\omega(S_n(\omega, \cdot) > n\varepsilon) = -c(\varepsilon), \quad \text{for } 0 < \varepsilon < \varepsilon_0 \text{ and a.e. } \omega.$$

We can also obtain CLT.

### Theorem (Central limit theorem)

If  $\Sigma^2 > 0$ , we have that

$$\lim_{n \rightarrow \infty} \int g(S_n(\omega, \cdot)/\sqrt{n}) d\mu_\omega = \int g dN(0, \Sigma^2),$$

for  $g$  continuous and bounded and a.e.  $\omega \in \Omega$ .

# Idea of the proof

We need to show that

$$\lim_{n \rightarrow \infty} \int e^{it \frac{S_n(\omega, \cdot)}{\sqrt{n}}} d\mu_\omega = e^{-\frac{t^2 \Sigma^2}{2}}, \quad \text{for a.e. } \omega \in \Omega.$$

This follows by proving that:

①

$$\lim_{n \rightarrow \infty} \int e^{it \frac{S_n(\omega, \cdot)}{\sqrt{n}}} d\mu_\omega = \lim_{n \rightarrow \infty} \prod_{j=0}^{n-1} \lambda_{\sigma^j \omega}^{\frac{it}{\sqrt{n}}},$$

where

$$\lambda_\omega^\theta = \int L_{\sigma^{-1}\omega}^\theta (v_{\sigma^{-1}\omega}^0 + \mathcal{W}^\theta(\sigma^{-1}\omega, \cdot)) dm =: H(\theta, \mathcal{W}^\theta)(\omega);$$

② by Taylor expansion of  $\theta \rightarrow H(\theta, W^\theta)$  around 0:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \log \lambda_{\sigma^j \omega}^{\frac{it}{\sqrt{n}}} = -\frac{t^2 \Sigma^2}{2}.$$

# Local central limit theorem

## Theorem (Local central limit theorem)

Assume that  $\Lambda(it) < 0$  for  $t \neq 0$ . Then, for a.e.  $\omega$  and every bounded interval  $J \subset \mathbb{R}$ , we have

$$\lim_{n \rightarrow \infty} \sup_{s \in \mathbb{R}} \left| \Sigma \sqrt{n} \mu_{\omega}(s + S_n g \in J) - \frac{1}{\sqrt{2\pi}} e^{-\frac{s^2}{2n\Sigma^2}} |J| \right| = 0,$$

