

**Baryon transition form factors at the pole**L. Tiator,<sup>1</sup> M. Döring,<sup>2,3</sup> R. L. Workman,<sup>2</sup> M. Hadžimehmedović,<sup>4</sup> H. Osmanović,<sup>4</sup> R. Omerović,<sup>4</sup> J. Stahov,<sup>4</sup> and A. Švarc<sup>5</sup><sup>1</sup>*Institut für Kernphysik, Universität Mainz, D-55099 Mainz, Germany*<sup>2</sup>*The George Washington University, Washington, DC 20052, USA*<sup>3</sup>*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA*<sup>4</sup>*University of Tuzla, Faculty of Natural Sciences and Mathematics, Univerzitetska 4, 75000 Tuzla, Bosnia and Herzegovina*<sup>5</sup>*Rudjer Bošković Institute, Bijenička Cesta 54, P.O. Box 180, 10002 Zagreb, Croatia*

(Received 11 June 2016; published 21 December 2016)

Electromagnetic resonance properties are uniquely defined at the pole and do not depend on the separation of the resonance from background or the decay channel. Photon-nucleon branching ratios are nowadays often quoted at the pole, and we generalize the considerations to the case of virtual photons. We derive and compare relations for nucleon to baryon transition form factors both for the Breit-Wigner and the pole positions. Using the MAID2007 and SAID SM08 partial wave analyses of pion electroproduction data, we compare the  $G_M$ ,  $G_E$ , and  $G_C$  form factors for the  $\Delta(1232)$  resonance excitation at the Breit-Wigner resonance and pole positions up to  $Q^2 = 5 \text{ GeV}^2$ . We also explore the  $E/M$  and  $S/M$  ratios as functions of  $Q^2$ . For pole and residue extraction, we apply the Laurent + Pietarinen method.

DOI: [10.1103/PhysRevC.94.065204](https://doi.org/10.1103/PhysRevC.94.065204)**I. INTRODUCTION**

As baryon resonance properties, evaluated at the pole position, are now beginning to supersede and replace quantities that historically have been determined using Breit-Wigner (BW) plus background parameterizations, we extend a recent study [1] of photodecay couplings at the pole to the regime of nonzero  $Q^2$ . The shift to pole-related quantities is reflected in the Review of Particle Properties (RPP) [2], with many pole values coming from the Bonn-Gatchina multichannel analyses [3]. Some plots of transition form factors at the BW position, as a function of photon virtuality, are now also available in the RPP.

Resonances are formally considered as poles in the  $S$  matrix, with a position independent of the reaction and with a factorizable residue for different reaction channels. The BW approach is more dependent on the formalism used to provide the associated background. While BW parameters have been fitted to electroproduction data in early analyses, more sophisticated approaches (such as dynamical models [4–6] and chiral effective theory) are not directly comparable to these values. In the chiral effective field theory calculation of Ref. [7], for example, complex form factor results were determined at the pole position. Comparisons to phenomenological fits providing real BW quantities were therefore *ad hoc*. Also, in chiral unitary resonance dynamics, there is no genuine resonance seed that would allow for the definition of a meaningful, purely real helicity coupling [8–10]. See also Refs. [11,12] for related results. Proper comparisons require pole parameters extracted from data with minimal model bias. We revisit these comparison below, once pole parameters have been determined.

While calculations at the pole are, in principle, well-defined and less model-dependent than the BW approach, the continuation of fit amplitudes to the pole is itself a possible source of error. With the aim to minimize model dependence of the pole extraction procedure, this uncertainty has motivated numerous studies involving speed plots, regularization methods, contour

integration [13–18], and the most recent Laurent series representations (L+P) based on separation of pole and regular parts, and using the conformal mapping variable to expand the regular part in power series [19]. Here we have extracted pole parameters with the L+P method from two energy-dependent (ED) partial wave analyses, MAID and SAID, which were fitted to the world database of pion electroproduction. The differences observed by this comparison will give an insight of the uncertainty of the pole form factors due to the differences in the MAID and SAID techniques and consequently also to the experimental database.

For the  $\Delta(1232)$  state, considerable attention [20–33] has been addressed to the  $Q^2$  evolution of amplitudes, as well as differences in the  $Q^2$  dependence of bare couplings, within models, and meson-cloud contributions. For a review, see, e.g., Ref. [34]. Transition form factors are now also determined in lattice QCD simulations [35,36]. Here, the quark masses are so large that the  $\Delta(1232)$  appears as a bound state, but it was realized in Ref. [37] that in future simulations, close to the physical point, the finite resonance width will complicate the extraction. Therefore, in Ref. [37] a method has been proposed to determine transition form factors at the pole. This stresses again the relevance of providing pole values for existing phenomenological analyses, which is the aim of this study.

The main focus of this paper is the  $\Delta(1232)$  virtual-photon decay amplitudes and related transition form factors. As the amplitudes themselves become infinite at the pole, we are interested in residues. The connection between multipole residues and the photodecay amplitudes has been clarified in a previous paper [1] and in the  $N$  and  $\Delta$  resonance minireviews of the 2012 and 2014 Particle Data Group listings [2].

Here we will compare BW and pole extractions, using the MAID2007 and SAID SM08 partial wave analyses of pion electroproduction data, utilizing the recent Laurent + Pietarinen (L+P) pole extraction method [38–42] which has proved to be a precise and very reliable tool for the determination of pole positions and residues.

The  $\Delta(1232)$   $E/M$  and  $S/M$  ratios have been studied for many decades. Interest in the  $E/M$  ratio, for real-photon interactions, was largely motivated by the fact that, in a simple nonrelativistic quark model, this ratio would be zero [43] and, thus, deviations from zero would require more complicated interactions. The measured value of this ratio was small [2],  $-2.5 \pm 0.5\%$ , but its precise value varied as photoproduction cross sections and beam-asymmetry  $\Sigma$  measurements became more precise. The prediction for this ratio, at very large  $Q^2$  from pQCD [44], has been more difficult to confirm. The ratio is predicted to become unity, whereas, at the real-photon point, it is small and negative. The variation of this ratio with  $Q^2$  has also changed significantly as electroproduction data have improved.

In Sec. II, we first give definitions of the standard BW quantities and then define the associated pole-valued results we will be considering. In Sec. III, we then give a brief overview of the BW and pole behaviors of the  $\Delta(1232)$  amplitudes, which are constrained by Watson's theorem. The L+P fit is described in Sec. IV and compared to results from a fit, generalized to nonzero  $Q^2$ , described in Ref. [1]. Finally, in Sec. IV, we summarize our findings and prospects for future work.

## II. BREIT-WIGNER VERSUS POLE QUANTITIES

The total cross section of pion electroproduction can be written as a semi-inclusive electron scattering cross section

$$\frac{d\sigma}{d\Omega_e dE_f} = \Gamma_V (\sigma_T + \varepsilon \sigma_L) \quad (1)$$

with the virtual-photon flux factor

$$\Gamma_V = \frac{\alpha}{2\pi^2} \frac{E_f}{E_i} \frac{\kappa_\ell}{\kappa_c} \frac{1}{1 - \varepsilon}, \quad (2)$$

where  $E_i$  and  $E_f$  are the initial and final electron energies in the laboratory frame, the virtual-photon polarization is  $\varepsilon$ , and the total transverse and longitudinal virtual-photon cross sections are

$$\sigma_T = \frac{1}{2} (\sigma_T^{1/2} + \sigma_T^{3/2}), \quad (3)$$

$$\sigma_T^h = 4\pi \frac{q}{\kappa_c} \sum_{\alpha(\ell, J, I)} (2J+1) |\mathcal{A}_\alpha^h|^2 C^2, \quad (4)$$

$$\sigma_L = 4\pi \frac{q}{\kappa_c} \frac{Q^2}{k^2} \sum_{\alpha(\ell, J, I)} (2J+1) |\mathcal{S}_\alpha^{1/2}|^2 C^2, \quad (5)$$

with  $q$  and  $k$  being the center-of-mass pion and photon momenta and  $\kappa_l = (W^2 - m_N^2)/2m_N$  and  $\kappa_c = (W^2 - m_N^2)/2W$  the so-called equivalent real photon energies in the laboratory and c.m. frames. The factor  $C$  is  $\sqrt{2/3}$  for isospin 3/2 and  $-\sqrt{3}$  for isospin 1/2. The helicity multipoles are given in terms of electric, magnetic, and longitudinal (timelike) multipoles:

$$\mathcal{A}_{\ell+}^{1/2} = -\frac{1}{2} [(\ell+2)E_{\ell+} + \ell M_{\ell+}], \quad (6)$$

$$\mathcal{A}_{\ell+}^{3/2} = \frac{1}{2} \sqrt{\ell(\ell+2)} [E_{\ell+} - M_{\ell+}], \quad (7)$$

$$\mathcal{S}_{\ell+}^{1/2} = -\frac{\ell+1}{\sqrt{2}} S_{\ell+}, \quad (8)$$

$$\mathcal{A}_{(\ell+)-}^{1/2} = -\frac{1}{2} [\ell E_{(\ell+)-} - (\ell+2)M_{(\ell+)-}], \quad (9)$$

$$\mathcal{A}_{(\ell+)-}^{3/2} = -\frac{1}{2} \sqrt{\ell(\ell+2)} [E_{(\ell+)-} + M_{(\ell+)-}], \quad (10)$$

$$\mathcal{S}_{(\ell+)-}^{1/2} = -\frac{\ell+1}{\sqrt{2}} S_{(\ell+)-}, \quad (11)$$

with  $J = \ell + 1/2$  for “+” multipoles and  $J = (\ell + 1) - 1/2$  for “-” multipoles, all having the same total spin  $J$ .

In analogy to photoproduction [1], we define the virtual-photon decay amplitudes

$$A_h^{BW} = C \sqrt{\frac{q_r}{\kappa_r} \frac{\pi(2J+1)M_r \Gamma_r^2}{m_N \Gamma_{\pi,r}}} \tilde{\mathcal{A}}_\alpha^h, \quad (12)$$

$$S_{1/2}^{BW} = C \sqrt{\frac{q_r}{\kappa_r} \frac{\pi(2J+1)M_r \Gamma_r^2}{m_N \Gamma_{\pi,r}}} \tilde{\mathcal{S}}_\alpha^{1/2}, \quad (13)$$

where  $\tilde{\mathcal{A}}_\alpha^{1/2}$ ,  $\tilde{\mathcal{A}}_\alpha^{3/2}$ , and  $\tilde{\mathcal{S}}_\alpha^{1/2}$  are the imaginary parts of the resonance amplitudes at the BW position  $W_r = M_r$ .

Similarly, we define the virtual-photon amplitudes at the pole position,

$$A_h^{\text{pole}} = C \sqrt{\frac{q_p}{\kappa_p} \frac{2\pi(2J+1)W_p}{m_N \text{Res}_{\pi N}}} \text{Res } \mathcal{A}_\alpha^h, \quad (14)$$

$$S_{1/2}^{\text{pole}} = C \sqrt{\frac{q_p}{\kappa_p} \frac{2\pi(2J+1)W_p}{m_N \text{Res}_{\pi N}}} \text{Res } \mathcal{S}_\alpha^{1/2}, \quad (15)$$

where the subscript  $p$  denotes quantities evaluated at the pole position.

The photon momenta,  $\kappa_r$  and  $\kappa_p$ , are photon equivalent energies and can be written as virtual-photon momenta at  $Q^2 = 0$ . The amplitudes,  $\mathcal{A}_\alpha^h$  and  $\mathcal{S}_\alpha^{1/2}$ , as well as the residues,  $A_h^{\text{pole}}$  and  $S_{1/2}^{\text{pole}}$ , are functions of the photon virtuality  $Q^2$ .

Through linear combinations, the helicity form factors can also be related to electric, magnetic, and charge form factors. These so-called Sachs form factors,  $G_E^*$ ,  $G_M^*$ , and  $G_C^*$ , are usually given in two different conventions by Ash [45] and by Jones and Scadron [46]. Both are related by a square-root factor,  $G_{JS}(Q^2) = G_{\text{Ash}}(Q^2) \times \sqrt{1 + Q^2/(m_N + M_r)^2}$ .

Here we will concentrate on the  $\Delta(1232)$  transition form factors and give the corresponding expressions. For transitions with different spin and parity, similar relations can be found; see [47,48].

For the  $\gamma N \Delta$  transition, Jones and Scadron [46] give the following relations between the total cross sections and the Sachs form factors:

$$\begin{aligned} \sigma_T + \varepsilon \sigma_L = & \frac{2\pi\alpha}{\Gamma_r m_N^2} \frac{k_r(Q^2)^2}{\kappa_r(1 + Q^2/(m_N + M_\Delta)^2)} \\ & \times \left( G_M^{*2}(Q^2) + 3G_E^{*2}(Q^2) + \varepsilon \frac{Q^2}{4M_\Delta^2} G_C^{*2}(Q^2) \right). \end{aligned} \quad (16)$$

In the convention of Ash, the Sachs form factors take the form

$$G_M^*(Q^2) = -c_\Delta(A_{1/2} + \sqrt{3}A_{3/2}), \quad (17)$$

$$G_E^*(Q^2) = c_\Delta\left(A_{1/2} - \frac{1}{\sqrt{3}}A_{3/2}\right), \quad (18)$$

$$G_C^*(Q^2) = \sqrt{2}c_\Delta \frac{2M_\Delta}{k_\Delta} S_{1/2}, \quad (19)$$

with  $W_r = M_\Delta$  and  $c_\Delta = [(m_N^3 \kappa_\Delta)/(4\pi\alpha M_\Delta k_\Delta^2)]^{1/2}$ , where  $k_\Delta = k_\Delta(Q^2) = k(M_\Delta, Q^2)$  and  $\kappa_\Delta = \kappa_c(M_\Delta) = k(M_\Delta, 0)$  are the virtual-photon momentum and the photon equivalent energy at resonance. Because the  $\Delta(1232)$  is very close to an ideal resonance, the real parts of the amplitudes vanish at  $W = M_\Delta$ , and the form factors can be directly expressed in terms of the imaginary parts of the corresponding multipoles at the (Breit-Wigner) resonance position,

$$G_M^*(Q^2) = b_\Delta(Q^2) \text{Im}\{M_{1+}^{(3/2)}(M_\Delta, Q^2)\}, \quad (20)$$

$$G_E^*(Q^2) = -b_\Delta(Q^2) \text{Im}\{E_{1+}^{(3/2)}(M_\Delta, Q^2)\}, \quad (21)$$

$$G_C^*(Q^2) = -b_\Delta(Q^2) \frac{2M_\Delta}{k_\Delta(Q^2)} \text{Im}\{S_{1+}^{(3/2)}(M_\Delta, Q^2)\}, \quad (22)$$

where  $b_\Delta(Q^2) = [(8m_N^2 q_\Delta \Gamma_\Delta)/(3\alpha k_\Delta^2(Q^2))]^{1/2}$ , with  $\Gamma_\Delta = 115$  MeV and  $q_\Delta = q(M_\Delta)$  being the pion momentum at resonance.

Similarly, we can define the Sachs form factors, using Ash's conventions, at the pole position:

$$G_M^{\text{pole}}(Q^2) = b_p(Q^2) \text{Res} M_{1+}^{(3/2)}(W_p, Q^2), \quad (23)$$

$$G_E^{\text{pole}}(Q^2) = -b_p(Q^2) \text{Res} E_{1+}^{(3/2)}(W_p, Q^2), \quad (24)$$

$$G_C^{\text{pole}}(Q^2) = -b_p(Q^2) \frac{2W_p}{k_p(Q^2)} \text{Res} S_{1+}^{(3/2)}(W_p, Q^2), \quad (25)$$

where  $b_p(Q^2) = [(16m_N^2 q_p)/(3\alpha k_p^2(Q^2) \text{Res}_{\pi N})]^{1/2}$ .

In the literature, the following ratios of multipoles have been defined:

$$R_{EM}(Q^2) = \frac{E_{1+}^{3/2}(Q^2)}{M_{1+}^{3/2}(Q^2)} = -\frac{G_E^*(Q^2)}{G_M^*(Q^2)}, \quad (26)$$

$$R_{SM}(Q^2) = \frac{S_{1+}^{3/2}(Q^2)}{M_{1+}^{3/2}(Q^2)} = -\frac{k_\Delta(Q^2)}{2M_\Delta} \frac{G_C^*(Q^2)}{G_M^*(Q^2)}. \quad (27)$$

The ratios at the pole position are given accordingly:

$$R_{EM}^{\text{pole}}(Q^2) = \frac{\text{Res} E_{1+}^{3/2}(Q^2)}{\text{Res} M_{1+}^{3/2}(Q^2)} = -\frac{G_E^{\text{pole}}(Q^2)}{G_M^{\text{pole}}(Q^2)}, \quad (28)$$

$$R_{SM}^{\text{pole}}(Q^2) = \frac{\text{Res} S_{1+}^{3/2}(Q^2)}{\text{Res} M_{1+}^{3/2}(Q^2)} = -\frac{k_p(Q^2)}{2W_p} \frac{G_C^{\text{pole}}(Q^2)}{G_M^{\text{pole}}(Q^2)}. \quad (29)$$

### III. AMPLITUDES AT THE BREIT-WIGNER POSITION AND AT THE POLE POSITION

In general, a pion photo- or electroproduction amplitude  $T_{\gamma,\pi}$ , or any multipole, can be written as a sum of resonance

and background contributions:

$$T_{\gamma,\pi}(W, Q^2) = T_{\gamma,\pi}^{\text{res}}(W, Q^2) + T_{\gamma,\pi}^{\text{bg}}(W, Q^2). \quad (30)$$

In order to obtain the BW amplitudes at the resonance position, the resonance part has to be modeled in BW form with energy-dependent partial widths for all possible decay channels and with energy-dependent phases in order to obey unitarity; see e.g. Ref. [49]. In general, this resonance-background separation is only possible in a model-dependent way [50]. Consequently, this also leads to some model dependence in the mass  $M_r$  and width  $\Gamma_r$  and for the amplitudes  $\mathcal{A}_\alpha^h$  and  $\mathcal{S}_\alpha^{1/2}$ . The only exception in the baryonic spectrum is the  $\Delta(1232)$ , which is purely elastic and therefore has a well-defined  $K$ -matrix pole,  $M_\Delta = 1232$  MeV, where the scattering phase  $\delta_{33}(M_\Delta) = \pi/2$ . For the  $\Delta(1232)$  this coincides with the Breit-Wigner resonance position. Due to Watson's theorem, also for pion photo- and electroproduction, the phase is exactly 90 degrees at resonance and the resonance-background separation is unique, as the background amplitude  $T_{\gamma,\pi}^{\text{bg}}(M_\Delta, Q^2) = 0$ .

This situation is very different at the pole position. Since only the resonance part of the total amplitude contains a pole, a model-dependent resonance-background separation is unnecessary. Therefore, the pole positions and also the residues are model independent. However, they can suffer from uncertainties arising from the analytical continuation of the amplitudes, determined from data on the real energy axis, into the lower part of the complex energy plane.

### IV. L+P EXPANSION

Finding the pole positions and the residues of baryon resonances can be a difficult task. Some first attempts, applied to  $\pi N$  scattering amplitudes, were carried out by Höhler [51] and Cutkosky [52]. The optimal method would be an analytic continuation into the complex plane, within a dynamical and analytical model, carefully considering all branch cuts from open channels, that generally produce many poles on different Riemann sheets, where only the pole closest to the physical axis is relevant. In many cases, however, this is not possible in practice, e.g., when partial wave amplitudes can only be evaluated on the physical axis. For these cases, Höhler proposed the speed-plot technique [51], which was later extended by the regularization method [53]. In the present study, we apply the Laurent-plus-Pietarinen (L+P) method based on separation of pole and regular part, and using the conformal mapping variable to expand the regular part in power series; the method which has proved to be most reliable and has been applied to different processes [38–42]. A major advantage of the L+P method is the fact that it is a global method, describing the amplitudes over a wide energy range, treating threshold effects in terms of physical and effective branch points. Most other methods use only partial wave information in a local region around the relevant resonance position.

In this study, we have adopted the multichannel Laurent-plus-Pietarinen method (MC L+P), developed in Ref. [42], to the single-channel case where the  $E$ ,  $M$ , and  $S$  multipoles must be treated simultaneously as they share the same pole with associated  $\pi N$  resonance coupling. One could therefore




FIG. 1. Figures showing the quality of the fit. From top to bottom we show all three multipoles at three different photon virtualities  $Q^2 = 0, 1, \text{ and } 5 \text{ GeV}^2$  for MAID2007 and SAID SM08 models. Black circles and brown squares are real and imaginary part of multipoles respectively. Blue solid lines are real parts and red dashed lines are imaginary parts of the L+P fit to the given model. Panels (a)–(c) show  $E_{1+}^{3/2}, M_{1+}^{3/2}, S_{1+}^{3/2}$  of the MAID solution and (d)–(f) the same for the SAID solution.

describe the method as a coupled-multipole Laurent-plus-Pietarinen (CM L+P) method:

$$T_{\gamma,\pi}^a(W) = \sum_{j=1}^{N_{\text{pole}}} \frac{x_j^a + i y_j^a}{W_j - W} + \sum_{k=0}^{K^a} c_k^a X^a(W)^k + \sum_{l=0}^{L^a} d_l^a Y^a(W)^l + \sum_{m=0}^{M^a} e_m^a Z^a(W)^m,$$

$$X^a(W) = \frac{\alpha^a - \sqrt{x_P^a - W}}{\alpha^a + \sqrt{x_P^a - W}}, \quad Y^a(W) = \frac{\beta^a - \sqrt{x_Q^a - W}}{\beta^a + \sqrt{x_Q^a - W}},$$

$$Z^a(W) = \frac{\gamma^a - \sqrt{x_R^a - W}}{\gamma^a + \sqrt{x_R^a - W}}, \quad D_{dp} = \sum_a^{\text{all}} D_{dp}^a,$$

$$D_{dp}^a = \frac{1}{2N_W^a} \sum_{i=1}^{N_W^a} \left\{ \left[ \frac{\text{Re } T_{\gamma,\pi}^a(W^{(i)}) - \text{Re } T_{\gamma,\pi}^{a,\text{exp}}(W^{(i)})}{\text{Err}_{i,a}^{\text{Re}}} \right]^2 + \left[ \frac{\text{Im } T_{\gamma,\pi}^a(W^{(i)}) - \text{Im } T_{\gamma,\pi}^{a,\text{exp}}(W^{(i)})}{\text{Err}_{i,a}^{\text{Im}}} \right]^2 \right\} + \mathcal{P}^a,$$

$$\text{Err}_{i,a}^{\text{Re}} = 0.05 \times \frac{\sum_{k=1}^{N_W^a} |\text{Re } T_{\gamma,\pi}^a(W^{(k)})|}{N_W^a} + 0.05 \times |\text{Re } T_{\gamma,\pi}^a(W^{(i)})|,$$

$$\text{Err}_{i,a}^{\text{Im}} = 0.05 \times \frac{\sum_{k=1}^{N_W^a} |\text{Im } T_{\gamma,\pi}^a(W^{(k)})|}{N_W^a} + 0.05 \times |\text{Im } T_{\gamma,\pi}^a(W^{(i)})|,$$

$$\mathcal{P}^a = \lambda_c^a \sum_{k=1}^{K^a} (c_k^a)^2 k^3 + \lambda_d^a \sum_{l=1}^{L^a} (d_l^a)^2 l^3 + \lambda_e^a \sum_{m=1}^{M^a} (e_m^a)^2 m^3;$$

$a$  multipole index ( $E_{l_{\pm}}, M_{l_{\pm}}, S_{l_{\pm}}$ )  
 $N_{\text{pole}}$  number of poles  
 $W_j, W \in \mathbb{C}$   
 $x_i^a, y_i^a, c_k^a, d_l^a, e_m^a, \alpha^a, \beta^a, \gamma^a, \dots \in \mathbb{R}$   
 $K^a, L^a, M^a, \dots \in \mathbb{N}$  number of Pietarinen coefficients in multipole  $a$   
 $N_W^a$  number of energies in multipole  $a$   
 $D_{dp}$  discrepancy function; the quantity to be minimized  
 $\mathcal{P}^a$  Pietarinen penalty function  
 $\lambda_c^a, \lambda_d^a, \lambda_e^a$  Pietarinen weighting factors  
 $\text{Err}_{i,a}^{\text{Re, Im}}$  minimization error of real and imaginary parts respectively  
 $x_P^a, x_Q^a, x_R^a \in \mathbb{R}$  (or  $\in \mathbb{C}$ ).

Here  $x_j^a + i y_j^a$  are the channel (multipole) residua which are left free for all three multipoles  $E$ ,  $M$ , and  $S$ , and  $W_j = M_j - i \frac{\Gamma_j}{2}$  are the pole positions of resonances  $j$ , which are kept fixed to the values obtained from L+P fits of the single  $M$  multipole obtained in the real photon case [40]. In addition, as we expect a similar analytic structure for all three multipoles, we have fixed the three branch points to have the same value:  $x_P^E = x_P^M = x_P^S, x_Q^E = x_Q^M = x_Q^S$ , and  $x_R^E = x_R^M = x_R^S$ .

The Pietarinen expansions formalize the simplest analytic form of functions having branch points at the Pietarinen-expansion parameters, and in this paper we use three Pietarinen expansions with expansion parameters  $x_P^a$ ,  $x_Q^a$ , and  $x_R^a$  to describe the analytic structure of the nonresonant background.

The first coefficient,  $x_P^a$ , is restricted to the unphysical region and effectively represents contributions from all singular parts below the  $\pi N$  threshold (all left-hand cuts including a circular cut). The second parameter,  $x_Q^a$ , is fixed to the pion threshold at  $W = 1.077$  GeV. The third branch-point,  $x_R^a$ , for MAID multipoles is left free and effectively accounts for all inelastic-channel openings in the physical domain. Its values are above  $\pi\pi N$  threshold. For SAID multipoles,  $x_R^a$  is fixed at the complex branch point  $\pi\Delta = (1.37 - i 0.04)$  GeV, and it effectively parametrizes all inelastic-channel openings in the physical domain and a resonance in the three-body intermediate state.

In the fitting procedure we have used two poles for  $P_{33}$  MAID amplitudes, and we used only one pole and a complex branch-point for  $P_{33}$  SAID amplitudes. However, as a complex branch-point describes a pole hidden in a two-body channel of a three-body intermediate state, SAID is described by two poles as well.

The L+P fit was compared to a method used to extract photodecay amplitudes at the pole in Ref. [1]. Residues were extracted from the SAID electroproduction multipoles for  $Q^2$  from 0.1 to 5.0 GeV<sup>2</sup>. Application to the SAID multipoles had the benefit of a known pole and cut structure and a narrower range of fit energies was required. The values obtained in this way, and those found using the L+P method, were not significantly different. This served as an independent test of

TABLE I. Magnetic, electric and charge transition form factors,  $E/M$ ,  $S/M$  ratios and photon decay amplitudes at  $Q^2 = 0$  for the Breit-Wigner and for the pole position compared between MAID and SAID solutions. The BW parameters used for the conversion factor are  $M_{\Delta} = 1232$  MeV and  $\Gamma_{\pi} = \Gamma_r = 115$  MeV, and the pole parameters are  $W_p = (1210 - 50i)$  MeV and  $\text{Res}_{\pi N} = 50 e^{-i47^\circ}$ . The form factors and ratios are dimensionless and the photon decay amplitudes are given in units of GeV<sup>-1/2</sup>. For the complex values at the pole position, we give absolute values with the same sign as for the BW values and a phase.

	MAID values			SAID values		
	BW	pole		BW	pole	
$G_M$	2.97	3.20	-4.7°	3.11	3.38	-3.5°
$G_E$	0.064	0.202	49°	0.051	0.181	54°
$G_C$	1.18	2.11	35°	1.30	2.31	34°
$R_{EM}$	-0.022	-0.063	53°	-0.016	-0.054	58°
$R_{SM}$	-0.042	-0.067	33°	-0.044	-0.069	30°
$A_{1/2}$	-0.131	-0.131	-20°	-0.139	-0.142	-18°
$A_{3/2}$	-0.247	-0.261	-7.7°	-0.258	-0.273	-6.8°
$S_{1/2}$	0.016	0.027	22°	0.018	0.030	21°




FIG. 2. Helicity transition form factors  $A_{1/2}$ ,  $A_{3/2}$ , and  $S_{1/2}$  compared at the BW and poles position. The black solid curves show the real BW results and the blue dashed and red dash-dotted lines show real and imaginary parts of the complex pole form factors. The left column shows the results with the Mainz-MAID analysis and the right column with the GWU-SAID analysis.

the L+P method applied to the electroproduction reaction. The L+P method was subsequently used exclusively to obtain results from both the SAID and MAID multipoles.

## V. RESULTS AND CONCLUSIONS

In the L+P analysis, the pion electroproduction multipoles  $E_{1+}^{(3/2)}(W, Q^2)$ ,  $M_{1+}^{(3/2)}(W, Q^2)$ , and  $S_{1+}^{(3/2)}(W, Q^2)$ , from MAID2007 and SAID SM08 were fitted from threshold up to 2 GeV in the center-of-mass energy. These multipoles, which are accessible via the MAID and SAID web pages [54], are displayed in Fig. 1.

For  $Q^2$  values near the real-photon point, we fitted amplitudes from  $Q^2 = 0$  to  $0.5 \text{ GeV}^2$  in increments of  $0.1 \text{ GeV}^2$ . We then examined  $Q^2$  values in increments of  $1 \text{ GeV}^2$  up to  $5 \text{ GeV}^2$ , finding this region to have a less rapid variation. At each value of  $Q^2$ , amplitudes were analyzed in steps of  $10 \text{ MeV}$ .

Representative fit results covering the  $\Delta(1232)$  energy range in Fig. 1 illustrate the very good fit quality and also

display the rapid fall off of these amplitudes with  $Q^2$ . Numerical results from the  $Q^2 = 0$  analyses are compiled in Table I.

In Fig. 2, we plot the associated helicity transition form factors,  $A_{1/2}$ ,  $A_{3/2}$ , and  $S_{1/2}$  as functions of  $Q^2$ . The  $A_{1/2}$  and  $A_{3/2}$  amplitudes, being dominated by the well-determined magnetic multipole, are very similar for the MAID and SAID analyses. The  $S_{1/2}$  variation in  $Q^2$  is qualitatively similar but differs in detail. It is interesting to note that, for the  $A_{1/2}$  and  $A_{3/2}$  amplitudes, the BW values and real parts of the pole quantities are nearly identical, particularly with increasing  $Q^2$ .

In Fig. 3, we compare the quantities  $G_M/G_D$ , and the  $E/M$  and  $S/M$  ratios as functions of  $Q^2$ , where  $G_D = (1 + Q^2/b^2)^{-2}$ , with  $b^2 = 0.71 (\text{GeV}/c)^2$ . Here also, the MAID and SAID results for  $G_M/G_D$  agree very closely, with only a small difference between the BW values and real parts of the pole quantities. This pole behavior has also been displayed, over a smaller  $Q^2$  range, in the analysis of Ref. [18]. The MAID and SAID BW results also agree well with the available single- $Q^2$  analyses of the  $E/M$  ratio. These plots give no




FIG. 3. Magnetic transition form factor in the notation of Ash divided by the dipole form factor  $G_M/G_D$  and  $E/M$  and  $S/M$  ratios compared at the BW and poles position. The black solid curves show the real BW results and the blue dashed and red dash-dotted lines show real and imaginary parts of the complex pole form factors. The left column shows the results with the Mainz-MAID analysis and the right column with the GWU-SAID analysis. The data points for  $G_M$  are from Refs. [58] (black circle), [59] (black squares), [60] (blue circles), [61] (green triangles) and [62] (red diamonds); for  $E/M$ : [58] (black circle), [63] (black diamonds), [64] (black triangle), [65,66] (black square), [67] (blue circle), [61] (green triangles) and [62] (red diamonds); and for  $S/M$ : [63] (black diamonds), [68] (black circle), [69] (black triangle), [65,66] (black square), [67] (blue circle), [61] (green triangles) and [62] (red diamonds).

indication of a crossover to positive  $E/M$  values, as expected from Ref. [44]. Previously, both the MAID (2003) [55] and SAID (2002) [56] fits had found indications for a crossover. This trend has disappeared with the incorporation of new and more precise measurements. The  $S/M$  ratios of the MAID and SAID analyses display the only qualitative difference in  $Q^2$  variation. Here also the BW and real parts of the pole behavior are similar, with the SAID (pole and BW) curves tending to approximately follow the behavior of the single- $Q^2$  fits, whereas the MAID trend is for a slower  $Q^2$  variation. We note that in the 2003 MAID analysis [55] the  $S/M$  ratio was found to have a more rapid  $Q^2$  variation, following the trend of existing single- $Q^2$  values.

For low values of  $Q^2$ , we can also compare to the expectations from chiral effective theory [7]. In Fig. 4, the MAID and SAID quantities from Fig. 3 are compared to

the predictions of Gail and Hemmert [7] over a restricted  $Q^2$  range. The range of applicability of their approach was estimated to about  $Q_{\max}^2 \approx 0.2 \text{ GeV}^2$ . Due to the lack of data at the pole position, single- $Q^2$  data extracted at the BW position were used to determine the parameters of their approach. The result is a qualitatively good agreement between the real parts of pole-valued quantities, especially for the dominant magnetic transition, where even the imaginary part is reasonably described. However, this is not the case for  $G_E$  and  $G_C$ . The real parts of these transitions are still in a moderate agreement, but the imaginary parts are off even by a different sign. This is not too surprising because the imaginary parts strongly depend on the parameters used for the pion loop integrals. A revised relativistic chiral perturbation theory calculation in the complex mass scheme [57] is in progress and may shed light on this issue.




FIG. 4. Magnetic, electric and charge transition form factors compared with the heavy baryon chiral effective field theory (HBCHEFT) of Gail and Hemmert [7] at low  $Q^2$ . The blue short-dashed and red dash-dotted lines show real and imaginary parts of the complex pole form factors obtained from MAID and SAID. The blue long-dashed and the red dotted lines are the real and imaginary parts of the HBCHEFT calculations.

#### ACKNOWLEDGMENTS

This work was supported in part by US Department of Energy Grant No. DE-SC0014133, by the Deutsche Forschungsgemeinschaft (SFB 1044), and by RFBR Grant

No. 13-02-00425. M.D. is supported by the National Science Foundation (CAREER Grant No. PHY-1452055 and PIF Grant No. 1415459) and by the US Department of Energy, Office of Science, Office of Nuclear Physics under Contract No. DE-AC05-06OR23177.

- 
- [1] R. L. Workman, L. Tiator, and A. Sarantsev, *Phys. Rev. C* **87**, 068201 (2013).  
 [2] K. A. Olive *et al.* (Particle Data Group), *Chin. Phys. C* **38**, 090001 (2014); J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).  
 [3] A. V. Anisovich, R. Beck, E. Klempt, V. A. Nikonov, A. V. Sarantsev, and U. Thoma, *Eur. Phys. J. A* **48**, 15 (2012).  
 [4] H. Kamano, S. X. Nakamura, T.-S. H. Lee and T. Sato, *Phys. Rev. C* **88**, 035209 (2013).  
 [5] D. Rönchen, M. Döring, H. Haberzettl, J. Haidenbauer, U.-G. Meißner, and K. Nakayama, *Eur. Phys. J. A* **51**, 70 (2015).  
 [6] D. Rönchen *et al.*, *Eur. Phys. J. A* **50**, 101 (2014); **51**, 63(E) (2015).  
 [7] T. A. Gail and T. R. Hemmert, *Eur. Phys. J. A* **28**, 91 (2006).  
 [8] D. Jido, M. Döring, and E. Oset, *Phys. Rev. C* **77**, 065207 (2008).  
 [9] M. Döring, D. Jido, and E. Oset, *Eur. Phys. J. A* **45**, 319 (2010).  
 [10] M. Döring, *Nucl. Phys. A* **786**, 164 (2007).  
 [11] V. Pascalutsa and M. Vanderhaeghen, *Phys. Rev. D* **73**, 034003 (2006).  
 [12] V. Bernard, T. R. Hemmert, and U. G. Meißner, *Phys. Lett. B* **622**, 141 (2005).  
 [13] M. Döring, C. Hanhart, F. Huang, S. Krewald, and U.-G. Meißner, *Nucl. Phys. A* **829**, 170 (2009).



- [14] S. Ceci, M. Döring, C. Hanhart, S. Krewald, U.-G. Meißner, and A. Svarc, *Phys. Rev. C* **84**, 015205 (2011).
- [15] P. Masjuan, in *The IX International Conference on Quark Confinement and the Hadron Spectrum—QCHS IX*, 2010, Madrid, edited by F. J. Llanes-Estrada and J. R. Peláez, AIP Conf. Proc. No. 1343 (AIP, New York, 2011), p. 334.
- [16] S. N. Yang, *Chin. J. Phys.* **49**, 1158 (2011).
- [17] L. Tiator, S. S. Kamalov, S. Ceci, G. Y. Chen, D. Drechsel, A. Svarc, and S. N. Yang, *Phys. Rev. C* **82**, 055203 (2010).
- [18] N. Suzuki, T. Sato, and T.-S. H. Lee, *Phys. Rev. C* **82**, 045206 (2010).
- [19] A. Svarc, M. Hadžimehmedovic, H. Osmanovic, and J. Stahov, [arXiv:1212.1295](https://arxiv.org/abs/1212.1295).
- [20] I. G. Aznauryan and V. D. Burkert, [arXiv:1603.06692](https://arxiv.org/abs/1603.06692).
- [21] G. Ramalho, M. T. Peña, J. Weil, H. van Hees, and U. Mosel, *Phys. Rev. D* **93**, 033004 (2016).
- [22] I. G. Aznauryan and V. D. Burkert, *Phys. Rev. C* **92**, 035211 (2015).
- [23] J. Segovia, I. C. Cloët, C. D. Roberts, and S. M. Schmidt, *Few-Body Syst.* **55**, 1185 (2014).
- [24] J. Segovia, C. Chen, I. C. Cloët, C. D. Roberts, S. M. Schmidt, and S. Wan, *Few-Body Syst.* **55**, 1 (2014).
- [25] M. Ronniger and B. C. Metsch, *Eur. Phys. J. A* **49**, 8 (2013).
- [26] H. Sanchis-Alepuz, R. Williams and R. Alkofer, *Phys. Rev. D* **87**, 096015 (2013).
- [27] G. Ramalho and M. T. Peña, *Phys. Rev. D* **85**, 113014 (2012).
- [28] E. Santopinto and M. M. Giannini, *Phys. Rev. C* **86**, 065202 (2012).
- [29] I. G. Aznauryan and V. D. Burkert, *Prog. Part. Nucl. Phys.* **67**, 1 (2012).
- [30] E. Santopinto, A. Vassallo, M. M. Giannini, and M. De Sanctis, *Phys. Rev. C* **82**, 065204 (2010).
- [31] B. Juliá-Díaz *et al.*, *Phys. Rev. C* **80**, 025207 (2009).
- [32] M. Ungaro *et al.*, *Phys. Rev. Lett.* **97**, 112003 (2006).
- [33] M. Fiolhais, B. Golli, and S. Sirca, *Phys. Lett. B* **373**, 229 (1996).
- [34] V. Pascalutsa, M. Vanderhaeghen, and S. N. Yang, *Phys. Rep.* **437**, 125 (2007).
- [35] C. Alexandrou, J. W. Negele, M. Petschlies, A. Strelchenko, and A. Tsapalis, *Phys. Rev. D* **88**, 031501 (2013).
- [36] C. Alexandrou, G. Koutsou, J. W. Negele, Y. Proestos, and A. Tsapalis, *Phys. Rev. D* **83**, 014501 (2011).
- [37] A. Agadjanov, V. Bernard, U.-G. Meißner, and A. Rusetsky, *Nucl. Phys. B* **886**, 1199 (2014).
- [38] A. Švarc, M. Hadžimehmedović, H. Osmanović, J. Stahov, L. Tiator, and R. L. Workman, *Phys. Rev. C* **88**, 035206 (2013).
- [39] A. Švarc, M. Hadžimehmedović, R. Omerović, H. Osmanović, and J. Stahov, *Phys. Rev. C* **89**, 045205 (2014).
- [40] A. Švarc, M. Hadžimehmedović, H. Osmanović, J. Stahov, L. Tiator, and R. L. Workman, *Phys. Rev. C* **89**, 065208 (2014).
- [41] A. Švarc, M. Hadžimehmedović, H. Osmanović, J. Stahov, and R. L. Workman, *Phys. Rev. C* **91**, 015207 (2015).
- [42] A. Švarc, M. Hadžimehmedović, H. Osmanović, J. Stahov, L. Tiator, and R. L. Workman, *Phys. Lett. B* **755**, 452 (2016).
- [43] C. Becchi and G. Morpurgo, *Phys. Lett.* **17**, 352 (1965).
- [44] C. E. Carlson, *Phys. Rev. D* **34**, 2704 (1986).
- [45] W. W. Ash *et al.*, *Phys. Lett. B* **24**, 165 (1967).
- [46] H. F. Jones and M. D. Scadron, *Ann. Phys. (NY)* **81**, 1 (1973).
- [47] R. C. E. Devenish, T. S. Eizenschitz, and J. G. Körner, *Phys. Rev. D* **14**, 3063 (1976).
- [48] L. Tiator, D. Drechsel, S. S. Kamalov, and M. Vanderhaeghen, *Eur. Phys. J. Spec. Top.* **198**, 141 (2011).
- [49] D. Drechsel, S. S. Kamalov, and L. Tiator, *Eur. Phys. J. A* **34**, 69 (2007).
- [50] M. Döring, C. Hanhart, F. Huang, S. Krewald, and U.-G. Meißner, *Phys. Lett. B* **681**, 26 (2009).
- [51] G. Höhler,  *$\pi N$  Newsletter* **9**, 1 (1993).
- [52] R. E. Cutkosky, C. P. Forsyth, R. E. Hendrick, and R. L. Kelly, *Phys. Rev. D* **20**, 2839 (1979).
- [53] S. Ceci, J. Stahov, A. Švarc, S. Watson, and B. Zauner, *Phys. Rev. D* **77**, 116007 (2008).
- [54] Amplitudes are available from the MAID site, <http://portal.kph.uni-mainz.de/MAID> and from the SAID site <http://gwdac.phys.gwu.edu>
- [55] L. Tiator, D. Drechsel, S. S. Kamalov, and S. N. Yang, *Eur. Phys. J. A* **17**, 357 (2003).
- [56] R. A. Arndt, W. Briscoe, I. Strakovsky, and R. Workman, in *Proceedings of the Workshop on the Physics of Excited Nucleons (NSTAR2002)*, Pittsburgh, 2002, edited by S. A. Dytman and E. S. Swanson (World Scientific, Singapore, 2003), p. 234.
- [57] M. Hilt, T. Bauer, S. Scherer, and L. Tiator (unpublished).
- [58] R. Beck *et al.*, *Phys. Rev. C* **61**, 035204 (2000).
- [59] W. Bartel *et al.*, *Phys. Lett. B* **28**, 148 (1968).
- [60] S. Stein *et al.*, *Phys. Rev. D* **12**, 1884 (1975).
- [61] V. V. Frolov *et al.*, *Phys. Rev. Lett.* **82**, 45 (1999).
- [62] I. G. Aznauryan *et al.* (CLAS Collaboration), *Phys. Rev. C* **80**, 055203 (2009).
- [63] S. Stave *et al.* (A1 Collaboration), *Phys. Rev. C* **78**, 025209 (2008).
- [64] C. Mertz *et al.*, *Phys. Rev. Lett.* **86**, 2963 (2001).
- [65] R. W. Gothe, *Prog. Part. Nucl. Phys.* **44**, 185 (2000).
- [66] T. Bantes, Ph.D. thesis, University of Bonn, 2003 (unpublished).
- [67] J. J. Kelly *et al.*, *Phys. Rev. Lett.* **95**, 102001 (2005); *Phys. Rev. C* **75**, 025201 (2007).
- [68] Th. Pospischil *et al.*, *Phys. Rev. Lett.* **86**, 2959 (2001).
- [69] D. Elsner *et al.*, *Eur. Phys. J. A* **27**, 91 (2006).